


- [Home](#)
- [Editorial Board](#)
- [Submit article](#)
- [Join as Reviewer/Editor](#)
- [List of Reviewer](#)
- [Most popular articles](#)
- [Free Articles](#)
- [Purchase Single Articles](#)
- [Archive](#)
- [Current Issue](#)
- [Recommend this journal to your library](#)
- [Advertiser](#)
- [Accepted Articles](#)
- [Search Articles](#)



Indian Journal of Pathology: Research and Practice

Volume 6, Issue 2 (Part-1), April - June 2017, Pages 298-302

Original Article

Study of Platelet Indices in Patients with Febrile Thrombocytopenia

Ramu R.* , Mahesh H. Karigoudar**, Yelikar B.R.***, Potekar R.M.**

*Assistant Professor, Department of Pathology, Basaveshwara Medical College, Hospital & Research Center, Chitradurga- 577501, Karnataka. **Professor ***Professor & Head, Department of Pathology, BLDE University™s Shri. B.M. Patil Medical College Ho

Choose an option to locate / access this Article:

Check if you have access through your login credentials.

DOI: <http://dx.doi.org/10.21088/ijprp.2278.148X.6217.23>

Abstract

Background: Platelet indices viz. Mean platelet volume (MPV), Platelet distribution width (PDW) and Platelet large cell ratio (P-LCR) are measured in the automated cell counter. These indices are well utilized for certain conditions like idiopathic thrombocytopenic purpura, aplastic anemia and other hemotological disorders to assess the prognosis.

 [Email Alerts](#)

 [Subscribe](#)

 [FAQ](#)

 [Contact Us](#)

Platelet count is decreased in various conditions including infectious and non-infectious disease processes. Infectious conditions like dengue fever, malaria, rickettsia, leptospirosis cause fever and variable degree of thrombocytopenia. Hence a detailed study of platelet indices is undertaken in patients having fever with thrombocytopenia. Materials & Methods: Present study was a prospective study. A total of 150 patients were included over a period of 1 year from Jan 2011 to Dec 2011. Detailed histories, physical and clinical examination of the patients were done to assess the etiology of febrile thrombocytopenia. Serological investigations for dengue fever, leptospirosis and rickettsia were done for confirmation. Results: Majority of the cases were of viral fever (72.6%), which includes dengue fever (10.6%), followed by malaria (15.3%), septicemia (4.6%), enteric fever (4%), leptospirosis (2%), rickettsia & brucellosis (0.6%). The platelet indices PDW (74.6%), MPV (86.6%), P-LCR (80.6%) showed less significant changes. Conclusion: Platelet indices are not changed significantly in febrile thrombocytopenic patients. Whenever patient with thrombocytopenia presents with changes in platelet indices, non infectious conditions should also be considered in differential diagnosis to manage the patients effectively for better clinical outcomes.

Keywords: Febrile Thrombocytopenia; Platelet Indices.

Corresponding Author : Ramu R.*

Information

[FAQ](#)
[Vacancy](#)
[Feedback](#)
[Term & Conditions](#)
[Privacy Policy](#)
[Cancellation and refund policy](#)

Resources

[Author/Editor](#)
[Librarian/Book Seller/Society](#)
[Institutions](#)
[Corporations](#)
[Conferences/Book Fairs](#)

Quick Links

[Rate list](#)
[Purchase single article](#)
[Order journal](#)
[Review article](#)
[Know status of your article](#)
[Submit article](#)

Contact Us

48/41-42, DSIDC, Pocket-II
Mayur Vihar Phase-I
Delhi - 110 091(India)
Phone: 91-11-22756995
Email : info@rfppl.co.in

