

Morphometric analysis of Infra orbital foramen position in south Indian skulls

Hussain Saheb Shaik¹, Muralidhar P Shepur², S.D Desai³, S.T Thomas⁴, GF Maavishettar⁵, Haseena S⁶

^{1, 2, 4 & 5} Department of Anatomy, JJM Medical College, Davangere, Karnataka- 577004, India

^{1, 3} Department of Anatomy, Shri BM Patil Medical College, Bijapur, Karnataka - 586 103, India

⁶ Department of physiology, Shri BM Patil Medical College, Bijapur, Karnataka - 586 103, India
anatomyshs@gmail.com

Corresponding author: Assistant Professor of Anatomy, JJMMC, Davangere -577004.

Email: anatomyshs@gmail.com

Abstract

The infra orbital foramen is an anatomical structure with an important location, through which the infra orbital vessels and nerve exits. The present study was conducted to determine the mean distance between the infra orbital foramen and infra orbital margin, mean distance between piriform aperture and infra orbital foramen, mean distance between the anterior nasal spine and infra orbital foramen on both side and as well as shape of foramen. Total 125 skulls were used for this study, the results were the mean distance between the infra orbital foramen and the infra orbital margin was 7.13 ± 1.78 mm, on right side it was 7.06 ± 1.81 mm and 7.20 ± 1.71 mm on left. The mean distance between the infra orbital foramen and the piriform aperture was 18.13 ± 1.85 mm, on right side it was 18.01 ± 2.03 mm and 18.25 ± 2.01 mm on left. The mean distance between infra orbital foramen and the anterior nasal spine on right was 33.97 ± 5.80 mm and left was 34.73 ± 4.70 mm. The knowledge of infra orbital foramen is very helpful for neurovascular surgeons.

Keywords: Infra orbital foramen, Infraorbital margin, Piriform aperture, Anterior nasal spine.

Introduction

The infra orbital foramen is situated bilaterally on the maxilar bone, lower to the edge of the orbital cavity, and it is bilateral. This foramen is directed inferior medial infra orbital nerve and vessels run through this foramen. The infra orbital nerve is sensitive and it lengthens the maxillary nerve, which crosses the infra orbital foramen and branches to feed the skin in the upper portion of the face, the maxillary sinus mucosa, the maxillary incisor, the canine and premolar teeth and the adjacent gum portion; the lower eyelid skin and conjunctiva, part of the nose, skin and mucosa of the upper lip (Apinhasmit *et al.*, 2006, Aziz *et al.*, 2000, Berge *et al.*, 2001)

Precise knowledge of the location of reference points in this area provides important data in local anesthesia and in maxillofacial and plastic surgical operations (Aziz *et al.*, 2000, Hwang *et al.*, 1999). There are several reference points on each wall with respect to infra orbital foramen, oral and maxillofacial surgery and local anesthesia (Du Tolt, 2003). Modern surgical procedures, anesthesia

(Zide, 1998), as well as acupuncture practice (Chonghuo, 1993), require more precise understanding of the surrounding anatomy (Bolini, 1990). The IOF is an important anatomical landmark that provides excellent analgesia for the closure of simple lacerations, biopsies, scar revisions, maxillofacial procedures, as well as various endoscopic and cosmetic cutaneous procedures.

Materials and Methods

125 dry adult human skulls constituted the material for the present study. The skulls belong to the Department of Anatomy, JJM Medical College, Davangere, Karnataka, India. Each was studied for the morphometric analysis of Infra orbital foramen position.

Results

Total 125 skulls were used for this study, the results were the mean distance between the infra orbital foramen and the infra orbital margin was 7.13 ± 1.78 mm, on right side it was 7.06 ± 1.81 mm and 7.20 ± 1.71 mm on left. The mean distance

between the infra orbital foramen and the piriform aperture was 18.13 ± 1.85 mm, on right side it was 18.01 ± 2.03 mm and 18.25 ± 2.01 mm on left. The mean distance between infra orbital foramen and the anterior nasal spine was 34.35 ± 5.25 mm, on right was 33.97 ± 5.80 mm and left was 34.73 ± 4.70 mm (Table 1). The transverse diameter of infra orbital foramen was 2.98 ± 0.69 mm and vertical diameter is 2.89 ± 0.71 mm (Table 2).

Table 1 :The mean distances between Infra orbital foramen and various reference points

Distances between	Right side	Left side	Total
IOF – IOM	7.06 ± 1.81 mm	7.20 ± 1.71 mm	7.13 ± 1.78 mm
IOF – PA	18.01 ± 2.03 mm	18.25 ± 2.01 mm	18.13 ± 1.85 mm
IOF – ANS	33.97 ± 5.80 mm	34.73 ± 4.70 mm	34.35 ± 5.25 mm

Table 2: Transverse Diameter and Verticle diameter of Infra orbital foramen

Measearment of IOF	
Transeverse Diameter	2.98 ± 0.69 mm
Verticle diameter	2.89 ± 0.71 mm

Discussion

The Infra orbital foramen is the way to the infraorbital nerve, vessels and the knowledge of its position is very useful to the professionals who manipulate the maxilar region like in acupuncture (Esper *et al.*, 1998), zygoma's fracture surgery (Du Tolt, 2003), practical of intra and extra oral anesthesia (Figun, 1994). The dentist and surgeons of head and neck have to know the exactly position of Infra orbital formen (Figun, 1994), because the anesthetic must be put on the foramen in order that it diffuses by the canal and causes the anterior superior alveolar nerve block and in consequence, the block of the branches that supply the central superior incisive teeth, lateral incisive and superior canine, ipsilateral to the blocked nerve (Zide, 1998).

Figun (1994) point out that, the topography of Infra orbital foramen presents unquestionable interesting on the anesthesia practice of alveolar anterior superior nerve and the infraorbital branch, because the foramen is an excellent reference point to intraoral functions and extra oral, 5 to 7mm inferior to the infra orbital margin. Dubrul (1991) reported about the variable distance between 6 to

8mm under the inferior margin of the orbit. Bergman *et al* (1988) describe a bigger variation, between 3 to 7mm inferior to the infra orbital margin. The distance between the Infra orbital foramen and midpoint of the inferior orbital margin has been reported to be from 4 mm to over 10 mm in several studies (Bolini 1993, Ochs, 1993, Chung *et al.*, 1995, Leo *et al.*, 1995).

Rontal *et al* (1979) found the distance between the Infra orbital foramen and midpoint of the inferior orbital fissure to be 24 mm. Karakas *et al.* (2002) pointed that the measurement between the Infra orbital foramen and midpoint of the inferior orbital margin was taken as 7 mm, Monod and Wanverts located an IOF on a line which is between the supraorbital incisure and the lip's comissure. According to the literature there are some standards of location of the foramen on the surface (Dubrul 1991), one of them is to check the line that joins the gengival margin, situated between the central and lateral incisive, medial to the fronto-zigomatic suture, a reference point easily recognized 4mm over a horizontal line that passes by the lateral angle of eyelid slit. Another classic conduct suggests situate the foramen on a vertical line that joins the supraorbital incisure to the mentonian foramen, 5 to 6mm under the orbital margin. Information on skull foramina size and symmetry (Del Sol *et al.*, 1989) is increasingly important because of the advancements in radiologic techniques such as magnetic resonance imaging and computed tomography. These methods are making difficult diagnoses of pathologic conditions of skull foramina possible (Berge, 2001).

The importance of the incidence and lateralization of the Infra orbital foramen is also evident in facial surgical procedures. The recognition of the presence of double or triple foramens is essential when the appropriate amount of anesthesia is applied, or it can be inappropriate. The study of the Infra orbital foramen is also basic to prevent the potential risk for iatrogenic injury during facial surgeries due to the presence of additional branches of the infra orbital nerve (Kazkayasi, 2003).

Previous studies show the relation between infra orbital foramen and the other anatomical structures, in studies with different purposes, what shows its importance as a repair point: distance between infra orbital foramen and an imaginary horizontal line in the piriform aperture base, inferior orbital fissure and the more inferior portion of the optical channel (Rontal, 1979) medium facial line on an imaginary line that passes through the supraorbital incisure; eyes pupil and second premolars (Molliex *et al.*, 1995) medium sagittal plane and supraorbital incisure (Chung, 1995) and finally related to the piriform aperture (Kazkayasi *et al.*, 2001).

References

1. Apinhasmit W, Chompoopong S, Methathrathip, D. et al (2006). Supraorbital notch/foramen, infraorbital foramen and Mental foramen in thais: anthropometric measurements and surgical Relevance. *Journal of the medical association of thailand.* 89(5), pp: 675-682.
2. Aziz SR, Marchena JM, Puran A (2000). Anatomic characteristics of the infra orbital foramen: a cadaver study. *Journal of oral maxillofacial surgery.* 58(9), pp: 992-996.
3. Berge JK, Bergman RA (2001). Variations in size and in symmetry of foramina of the human skull. *Clinical anatomy.* 14(6), pp: 406-413.
4. Hwang K, Baik S (1999). Surgical anatomy of Korean adults. *Journal of Craniofac. Surg.*, 10:129-34.
5. Du tolt DF, Nortje C (2003). The maxillae: integrated and Applied anatomy relevant to dentistry. *Sadj*, 58(8), 325-30.
6. Zide B, and Swift R (1998). How to block and tackle the face. *Plast. Reconstr. Surg.*, 101:2018.
7. Chonghuo T (1993). *Tratado de medicina chinesa.* Sao paulo, Roca.
8. Bolini P, Del sol M (1990). Considerações anatomicas sobre eo sulco infra-orbital. *Rev. Bras. Oftal.* 49(2):113-6.
9. Esper R, Yara J, Yamamura Y, Cricenti SV (1998). Relações Anatomicas do ponto de acupuntura e-2 (sibai) localizado No forame infra-orbital. *Rev. Paul. Acupunt.* 4(1):19-22.
10. Figun, ME, Garino R (1994). *Anatomia odontologica funcional E aplicada.* Sao paulo, panamericana.
11. Dubrul EL (1991) *Sicher and Dubrul oral anatomy.* 8th ed. *Ishiyaku euroamerica publishers.*
12. Bergman RA, Thompson SA, Afifi AK, Saadeh FA (1988) *Compendium of human anatomic variation: catalog, Atlas and world literature.* Baltimore. Urban & Schwarzenberg.
13. Ochs MW, Buckley M (1993). Anatomy of the orbit. *Oral Maxillofac. Surg. Clin. North. Am.* 5:419.
14. Leo JT, Cassell M, Bergman R (1995). Variations in human Infraorbital nerve, canal and foramen. *Ann. Anat.*, 177: 93.
15. Chung MS, Kim HJ, Kang HS, Chung IH (1995). Locational relationship of the supraorbital notch or foramen and infraorbital and mental foramina in Koreans. *Acta anat.* 154:162-6.
16. Rontal E, Rontal M, Guilford F T (1979). Surgical anatomy of the orbit. *Ann. Otol.* 88:382-6.
17. Karakas P, Bozkir M.G, Oguz O (2002). Morphometric measurements from various reference points in the orbit of male caucasians. *Surg. Radiol. Anat.*, 24(6), 358-62.
18. Kazkayasi M, Ergin A, Ersoy M *et al* (2003). A Microscopic anatomy of the infra orbital canal, nerve, and foramen. *Otolaryngology - Head & neck surgery.* 129, (6), 692-697.
19. Molliex S, Navez M and Baylot D (1996). Regional anesthesia for outpatient nasal surgery. *British journal of anaesthesia.* 76, (1) 151-153.