

Elongated Styloid Process (Eagle's Syndrome) : A Case Report

Kiran. V. Padeyappanavar, Ameer Khusru. M. Kazi. & S. D. Desai
 Belgaum Institute of Medical Sciences, Civil Hospital Compound,
 Belgaum - 590001, Karnataka

Abstract : During routine osteology discussion classes for under graduate students, it was observed in 3 of the skulls that their styloid processes were elongated on both the sides approximately measuring more than 5 cm in length .

Key words: Styloid process , skull

Introduction :

Elongated styloid process or calcified stylohyoid ligament can cause recurrent throat pain along with foreign body sensation , dysphagia or facial pain . Additional symptoms may include neck or throat pain with radiation to ipsilateral ear.

In adults the styloid process is approximately 2.5 cm in length and its tip is located between the external and internal carotid arteries , just lateral to the tonsillar fossa.


Material And Methods :

During the routine course of osteology discussion with undergraduate students it was observed in three dried human skulls approximately of 40 to 50 years old , their styloid processes were elongated.

Observation And Results :

The length of the styloid processes were measured with the help of a measuring tape.

Skull	Length in Cm.	
	Right	Left
A	4.5	5.0
B	4.0	5.5
C	5.2	5.0



Photographs of skulls showing elongated styloid processes.

Gross Anatomy :

The styloid process is a slender, elongated, cylindrical bony projection that lies anteromedial to the mastoid process . It normally varies in length from 2 cm. to 3 cm, and a styloid process longer than 3 cms is found in 4 to 7% of the population. The styloid process has attachments to three muscles and two ligaments . The stylopharyngeus, stylohyoid and styloglossus muscles originate here. The Facial nerve emerges from the stylomastoid foramen posteriorly. The stylohyoid ligament extends from the styloid process to the lesser cornu of the hyoid bone.

Embryology :

The styloid process , stylohyoid ligament and lesser cornu of the hyoid bone are derived from Reichert's cartilage , which arises from the second branchial arch. The cause of elongation of the styloid process has not been fully elucidated . Several theories have been proposed .

Correspondence to :-
Dr. Kiran. V. Padeyappanavar
 Belgaum Institute of Medical Sciences,
 Civil Hospital Compound,
 Belgaum - 590001, Karnataka.
 Mob. : 9886317512
 E-mail : dr.kiranvp@gmail.com

1. Congenital elongation of the process due to persistence of a cartilaginous anlage in the stylohyale.
2. Calcification of the stylohyoid ligament giving the appearance of an elongated styloid process.
3. Growth of Osseous tissue at the insertion of the stylohyoid ligament .

The third theory is based on histologic evidence of metaplastic changes to the subperiosteal cells in the vicinity of the ligaments insertion. Regardless of the pathophysiology of elongation , the result is a rigid , abnormally long structure that can cause pain or discomfort by one or several mechanisms.

Discussion :

Elongated styloid process or calcified stylohyoid ligament can cause recurrent throat pain along with foreign body sensation , dysphagia or facial pain.

Elongated styloid process is a diagnosis that should be considered in the evaluation of recurrent neck , throat or facial pain and dysphagia .

Additional Symptoms may include neck or throat pain with radiation to the ipsilateral ear. In adults the styloid process is approximately 2 to 3 cms in length and its tip is located between external and internal carotid arteries just lateral to the tonsillar fossa. It may develop inflammatory changes or impinge on the adjacent arteries , on sensory nerve endings leading to the symptoms described . Diagnosis can usually be made on physical examination by digital palpation of the styloid process in the tonsillar fossa which exacerbates the pain . In addition relief of symptoms with injection of an anaesthetic solution in to the tonsillar fossa is highly suggestive of this diagnosis . Radiographic work up should include antero-posterior and lateral skull films. The treatment of Eagles syndrome is primarily surgical. The styloid process can be shortened through an intraoral or external approach .

The first report of the Anatomic pathology was by Marchetti of Padua (1952) who described the ossification of the stylohyoid ligament.

The pain syndrome was recognize by Weinlecher (1872) surgically excise the styloid process to treat a symptomatic patient .

Other isolated cases were reported until Eagle described the syndrome in (1937) . Eagle divided the syndrome into two categories . He described the classic syndrome as persistent pain in the pharynx , aggravated by swallowing with the pain frequently referred to the ear on the side of the elongated styloid process. He also notice increased salivation , hesitancy , difficulty in swallowing , gagging and a foreign body sensation .

Fritz (1940) reported that only 11 of 43 patients with the syndrome in his series has had tonsillectomy . Both Eagle and Fritz reported that their patients were completely relieved of symptoms by the intraoral shortening of the styloid process.

Harma (1967) noted that bilateral elongation occurred in 50 % of the patients but only half of them had bilateral symptoms .

Steinmann (1968) reported the syndrome in 30 patients , 26 of whom did not have an elongated styloid process.

Keur et al (1986) assessed 1135 edentulous patients clinically and radio graphically to determine the relationship between the elongated styloid process and four symptoms frequently encountered in patients with Eagles syndrome.

Conclusion :

Eagle's syndrome though it is a rare entity it is largely under diagnosed . A thorough clinical and radiological examination will reveal impending insult. Proper diagnosis can definitely be of immense help to rationalize the line of management and the ultimate clinical out come.

References :

1. Eagle WW: Elongated styloid process . Arch Otolaryngol 1937; 25:584-587.
2. Eagle WW: Elongated styloid process: Symptoms and treatment . Arch Otolaryngol 1958; 64:172-176.
3. Fritz M: Elongated styloid process : cause of obscure throat symptoms . Arch Otolaryngol 1940; 31:911.
4. Hama R: Stylalgia . Acta otolaryngol Suppl (Stockh) 1967; 224: 149-155.
5. Steinmann EP : Styloid syndrome in absence of an elongated process. Acta Otolaryngol (Stockh) 1968; 66:347-356.
6. Keur JJ ,Campbell JPS, McCarthy JF, et al: The clinical significance of the elongated styloid process . Oral Surg Oral Med Oral pathol 1986; 61:399-404.
7. Holinshead WH: Anatomy for Surgeons .The Head and Neck. Philadelphia, JB Lippincott Co, 1982.
8. Steinmann EP : A new light on the pathogenesis of the styloid syndrome . Arch Otolaryngol 1970; 91:171-174.
9. Balasubramanian S: The ossification of the stylohyid ligament and its relation to facial pain. Br Dent J 1964; 116:108-111.
10. Strauss M, Yohar Z, Laurian N; Elongated styloid process syndrome : intraoral vs.external approach for styloid surgery . Laryngoscope 1985; 95:976-979.

**Proceedings
of
55th NATCON of ASI 07
held at LTMMC,SION (MUMBAI)**

**Office Bearers of A.S.I.
Year 2008**

Dr. K.K.Krishnamma (Kochi)
President
Dr. G.B. Rairam (Bagalkot)
Vice President
Dr. Pushpa Dhar (New Delhi)
Vice President
Dr. G.S.Longia
General Secretary
Dr. A.S. Mangalgiri (Bhopal)
Joint Secretary
Dr.Ajai Kumar Srivastava (Lucknow)
Treasurer
Dr. Punita Manik (Lucknow)
Joint Treasurer
Dr. S. L. Jethani (Dehradun)
Editor
Dr. Namita Mehrotra (Dehradun)
Joint Editor

**Executive Committee Members
For the year 2008:**

Dr. P.K. Verma (Mumbai)
Dr. Narga Nair (Manipal)
Dr. Royana Singh (Varanasi)
Dr.Keshaw Kumar (Allahabad)
Dr. B.R. Zambare (Ahmed Nagar)
Dr.Satyam Khare (Meerut)
Dr. Dharam Singh(Agra)
Dr. Vinod Kumar (Patna)
Dr. Parthapratim Pradhan (Kolkata)
Dr. D.S. Joshi (Miraj)
Dr. N.Hima Bindu (Moinabad)
Dr. Shova Rawlani (Wardha)
Dr. M.A.Doshi (Karad)
Dr. R.Chitra (Vijayawada)
Dr. B.Narasinga Rao