

Comparison of Cardiopulmonary Fitness among Obese and Non-Obese School Children

GAURAV ARORA¹, JYOTI P KHODNAPUR², SATISH G PATIL³, SUMANGALA M PATIL⁴

ABSTRACT

Introduction: Childhood obesity is now rising as a significant health problem. In India, some studies showed a rising trend in the prevalence of overweight and obesity in children. This shocking rise in childhood obesity often accounts for increased intake of high calorie foods and decreased physical activity. In addition, non communicable diseases like diabetes mellitus, cardiovascular diseases and cancer are more common in obesity.

Aim: The present study aimed to assess the relationship between cardiopulmonary fitness and Body Mass Index (BMI) in school going adolescents between 12 to 16 years.

Materials and Methods: This study was a cross-sectional observational study conducted from March 2018 to March 2019. The study included 60 healthy students (between 12 to 16 years) full sample size divided into two groups and four subgroups. After the general physical examination and history taking, participants' selection made in line with pre-defined inclusion and exclusion criteria. Anthropometric and body composition parameters were recorded. Using the modified Harvard's Step test, recorded cardiopulmonary fitness parameters like Physical

Fitness Index (PFI) and Maximal oxygen consumption (VO_2 max) of each subject and calculated by applying the concerned formulae. All statistical analysis has been done by using Statistical Package for the Social Sciences (SPSS) software version 16.0.

Results: Total 60 apparently healthy students including both male and female students. Mean age was 14.93 ± 0.96 years in Group I (normal weight boys, $n=15$) and 14.47 ± 1.41 years in Group II (overweight/obese boys, $n=15$). Also mean age was 14.93 ± 0.59 years in Group III (normal weight girls, $n=15$) and 15.27 ± 0.59 years in Group IV (overweight/obese girls, $n=15$). Pearson correlation showed a significant negative correlation between BMI and PFI ($r = -0.504$, p -value < 0.001) also between BMI and VO_2 max ($r = -0.459$, p -value < 0.001). This observation was explained by their sedentary activities (television viewing and video/computer time).

Conclusion: We all should take appropriate measures to enhance cardiopulmonary fitness among school children who are the wealth of the country at the school and community levels.

Keywords: Adolescents, Exercise, Physical fitness index, Obesity, Overweight

INTRODUCTION

In 2016, according to a study, more than 1.9 billion adults were overweight. Out of these, over 650 million were obese. Thirty-eight million children under the age of 5 were overweight or obese in 2019. Over 340 million children and adolescents aged 5-19 were overweight or obese in 2016 [1].

Developing countries face a double burden of disease, i.e., non communicable diseases (obesity and associated conditions, DM and Cardiovascular diseases) and infectious diseases, which impose a considerable and rapidly growing burden on low and middle income countries [2]. Physical fitness is the prime criterion for survival, to achieve any goal, and to lead a healthy life. Physical fitness indicates all the systems' functional status like skeletomuscular, cardio-respiratory, endocrine and metabolic [3].

Cardiopulmonary fitness is now considered an important health marker and a predictor of cardiovascular disease morbidity and mortality [4]. Even though physical fitness is genetically determined, it can also be influenced by environmental factors like sedentary lifestyle, reduced physical activity, and dietary habits [5]. Obesity is defined as abnormal or excessive fat accumulation that may impair health. WHO defines obesity as BMI > 30 kg/ m^2 [6]. Obesity is considered a major risk factor for coronary heart diseases, hypertension, diabetes mellitus, cerebrovascular accidents, depression, anxiety, Childhood and adolescence are crucial periods of life since dramatic physiological and psychological changes occur at these ages. Likewise, lifestyle and healthy/unhealthy behaviours are established during these years, influencing adult behaviour and future health status [7].

Previous studies have indicated that the frequency, duration, and intensity with which physical activity is crucial to an individual's overall health status suggest that a threshold must be maintained to produce positive health effects [8,9]. Maximal oxygen uptake (VO_2 max) is a standard measure of physical fitness levels, with higher values attained as fitness level improves, and is an essential indicator of successful physical activity interventions [10]. It's also important to note that most of the time dedicated to physical activity must be aerobic. Immediate measures must be taken to alleviate physical inactivity because it has been shown that: physical activity patterns track from childhood into adulthood, increasing physical activity modifies chronic disease (which also follow into adulthood) risk factor and prognosis, physical activity improves mental health as indicated by the lowered risk of development of anxiety, depression, in them. Increasing physical activity among children and adolescents is difficult as behaviour is influenced by physiological (age, gender, and ethnicity), psychological, sociocultural, and environmental factors [11].

Psychological determinants include self-perception of physical or sporting competence and personal benefits, possession of a positive mindset toward physical activity, level of engagement, and commitment. Barriers to physical activity, such as lack of time or feeling tired, are negatively associated with physical activity. Sociocultural influences include support for and participation in peers and siblings' physical activity, parental level of physical activity [12].

Since a child carries what he has learnt in school home, school is one of the cheapest and most efficient ways of health

communication. Most of the developing countries have a growing population with a substantial proportion at a young age, imparting timely health education regarding physical fitness will reduce the future disease burden of non communicable disease and be communicated across to the next generations [13].

Children with higher cardiorespiratory fitness levels are primarily not associated with cardiovascular disease and myocardial infarction in adulthood. Therefore, precise, proper, and clinically helpful quantification of cardiorespiratory fitness is essential for identifying low cardiorespiratory fitness levels in children [14-16].

The school setting is the bestsuited environment for population-based physical activity interventions as it plays a significant role in their development. Given this, the present study was designed to compare cardiopulmonary fitness among overweight/obese and normal weight school children. Hence, the hypothesis of this study predicted that BMI will affect the cardiopulmonary fitness. This study may provide school guidelines to implement health promotion policies and physical activity programs for those suffering from obesity and low cardiopulmonary fitness.

MATERIALS AND METHODS

A cross-sectional observational study conducted from March 2018 to March 2019 among school children in the age group of 12-16 years of Banjara high school in Vijayapur, Karnataka, India. The study obtained Institutional Ethical Certificate by the Institutional Ethical Committee (IEC Ref No-259/2017-18 dated March 27, 2018) Shri BM Patil Medical College, Hospital and Research Centre, BLDE (Deemed to be University).

Sample size calculation: Mean VO_2 max (mL/kg/min) by bicycle ergometry is 20.3 and the Standard Deviation (SD) is 5.5. the calculated sample size when the allowable error $L = \pm 2$ is:

$$n = \frac{4\sigma^2}{L^2} = \frac{4(5.5)^2}{(\pm 2)^2} \cong 30$$

Hence, a total of 60 subjects, including 30 overweight and obese, 30 normal weight school children are taken for the study. As all the distributions merged into a normal distribution, sample size, i.e., 30, was enough because inference that can draw based on 30 observations will more or less remain the same despite an increase in the sample size [5,17]. The participants were screened ($n=108$) in the school premises, and volunteers enrolled. Screening and recording for subjects were done in January 2020. Those subjects who were healthy were selected ($n=60$) for the study after a thorough examination based on inclusion and exclusion criteria.

Inclusion criteria: Total 60 Healthy school children of which 30 were overweight/obese (BMI was 23 kg/m² and above) [18] and 30 normal weight (BMI: 18.5-22.9 kg/m²) between 12-16 years of age were included in the study. The apparent health status of the subject was determined through clinical examination and history taking.

Exclusion criteria: Subjects with any congenital heart disease, endocrine disorders, history of chronic diseases, respiratory tract infection, allergy, cardiac or respiratory disease and congenital anomalies were excluded from the study.

The entire sample size was divided into 4 groups;

- Group I: normal weight ($n=15$; BMI: 18.5-22.9 kg/m²) boys,
- Group II: overweight/obese ($n=15$; BMI: 23 kg/m² and above) boys,
- Group III: normal weight ($n=15$; BMI: 18.5-22.9 kg/m²) girls,
- Group IV: overweight/ obese ($n=15$; BMI: 23 kg/m² and above) girls.

Each subject will be explained about the purpose and procedure to be adopted in the study. After taking consent from the principal, we first ruled out the exclusion criteria in the subjects by history. A thorough physical examination was done, followed by the recording of the anthropological and physiological parameters.

Anthropometric Parameters

Anthropometric parameters like height in centimetres (cm), weight in kilograms (kg), BMI in kilograms per square meter (kg/m²) and BSA in square meter (m²), Waist Circumference in centimetres (WC), Hip Circumference in centimetres (HC) and physiological parameters like pulse rate in (beats/min), systolic blood pressure in millimetre of mercury (mmHg), diastolic blood pressure (mmHg), pulse pressure (mmHg) and mean arterial pressure (mmHg) were recorded by using standard procedures [19-22]. WHR was calculated by dividing WC by HC [23,24].

Body Fat parameters: Body fat percentage (BF%), Fat Mass (FM), Fat Free Mass (FFM), Fat Mass Index (FMI) and Muscle Mass (MM) were recorded [25,26].

Cardiopulmonary Fitness Parameters

The test is done on the modified Harvard steps of 33 cm height and

1. Physical Fitness Index (PFI in %) [27]: The test is done on the modified Harvard steps of 33 cm height and it is calculated using following formula-

$$PFI = \frac{\text{Duration of exercise in sec}}{2 (\text{Sum of pulse counts during recovery})} \times 100$$

Procedure: The subjects were advised to step up on the modified Harvard steps of 33 cm height once every two seconds for 1minute. A total of 60 steps in total. The subjects were made to sit and the pulse rate was recorded immediately after the procedure and then at 1, 3, 5-minute intervals respectively, pulse rate is recorded as

- (a) Pulse Rate 0 (PR0) – immediately after exercise
- (a) Pulse Rate 1 (PR1) – 1 min after exercise
- (b) Pulse Rate 2 (PR2) – 3 min after exercise
- (c) Pulse Rate 3 (PR3) – 5 min after exercise

2. Maximal Aerobic Power (VO_2 Max) [28]: Pulse rate recovery was used as a proxy for VO_2 max. It is obtained by using the formula: $111.33 - (0.42 \times P \text{ max})$ [29]

P max: maximum pulse rate/min recorded immediately after 60 sec of Harvard's Step Test exercise.

STATISTICAL ANALYSIS

All statistical analysis has been done by using Statistical Package for the Social Sciences (SPSS) software version 16.0. All values are presented as Mean \pm SD and Standard Error (SE). A comparison of mean values of parameters between the groups has been done by using an unpaired t-test. The p-value ≤ 0.05 is taken as significant. The data has been expressed in the form of tables and graphs. Data were normally distributed. To find the correlation between the variables, the Pearson correlation coefficient (parametric) applied.

RESULTS

Among 60 Healthy school children, 30 (15 boys and 15 girls) overweight/obese (BMI: 23 kg/m² and above), 30 (15 boys and 15 girls) normal weight (BMI: 18.5-22.9 kg/m²) between 12-16 years of age showed significant high values of weight, BMI (p-value<0.001), WC (p-value<0.001), HC (p-value<0.001), WHR (p-value<0.001), BF% (p-value<0.001), FM (p-value<0.001), FFM (p-value=0.01), MM (p-value=0.01) and significant low values of PFI (p-value=0.03) and VO_2 max (p-value=0.03) among Group II (overweight/obese boys) compared to Group I (normal weight boys) [Table/Fig-1].

The study results also showed significant high values of weight, BMI (p=0.054), PR (p-value=0.025), PP (p-value=0.02), MAP (p-value=0.021), WC (p-value<0.001), HC (p-value=0.003), WHR (p-value=0.012), BF% (p-value<0.001), FM (p-value<0.001), FFM (p-value=0.014), MM (p-value=0.014) and significant low values of PFI (p-value<0.001) and VO_2 max (p-value<0.001) among Group IV (overweight/obese girls) compared to Group III (normal weight girls) [Table/Fig-2].

Parameters	Group I		Group II		t-values	p-values
	Mean±SD	SE	Mean±SD	SE		
Age(years)	14.93±0.96	0.25	14.47±1.41	0.36	1.060	0.3
Height (cm)	153.9±6.49	1.99	151.8±4.8	2.26	4.431	0.135
Weight (Kg)	48.1±5.27	1.27	57.83±6.67	2.35	2.017	<0.001
BMI (kg/m ²)	20.27±1.4	0.27	25.15±1.71	0.59	8.535	<0.001
BSA (m ²)	1.43±0.1	0.03	1.59±0.2	0.04	2	0.05*
PR (beats per minutes)	93.93±14.7	2.85	93.66±11.98	2.37	0.054	0.563
SBP (mmHg)	121.07±14.51	2.96	127.07±11.83	2.81	1.241	0.225
DBP (mmHg)	68.93±8.24	2.32	72.13±7.15	1.73	1.136	0.266
PP (mmHg)	52.13±6.32	3.06	56.33±7.12	3.78	0.908	0.372
MAP (mmHg)	86.31±8.3	2.10	90.9±6.4	1.75	1.695	0.101
WC (cm)	74.26±3.95	1.02	94.27±6.1	1.57	-10.7	<0.001
HC (cm)	88.33±4.42	1.14	103.93±7.9	2.04	-6.68	<0.001
WHR	0.84±0.05	0.01	0.91±0.06	0.02	-3.29	<0.001
RR (cpm)	18.4±1.88	0.49	18.67±2.38	0.61	-0.34	0.74
BF%	17.63±1.79	0.46	26.59±3.64	0.94	-8.55	<0.001
FM (kg)	9.36±1.27	0.33	17.96±4.16	1.07	-7.65	<0.001
FFM (kg)	43.77±4.17	1.08	49.11±6.14	1.59	-2.79	0.01**
MM (kg)	21.89±2.09	0.54	24.56±3.07	0.79	-2.79	0.01**
PFI (%)	49.19±3.51	0.91	46.56±2.6	0.67	2.327	0.03*
VO ₂ max (mL/kg/min)	57.35±3.3	0.85	54.52±3.4	0.88	2.315	0.03*


[Table/Fig-1]: Shows unpaired t-test results of anthropometric and physiological characteristics between Group I (normal weight boys) and Group II (overweight/obese boys). Data are Mean±SD and SE; Values in the final column represent results of unpaired t-test among two groups; Significant difference between groups showed at p<0.05 level; BMI: Body mass index; BSA: Body surface area; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; PP: Pulse pressure; MAP: Mean arterial pressure; WC: Waist circumference; HC: Hip circumference; PR: Pulse rate; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; WHR: Waist hip ratio; BF%: Body fat percentage; FM: Fat mass; FFM: Fat-free mass; FMI: Fat mass index; MM: Muscle mass; PFI: Physical fitness index and VO₂max: Maximal oxygen consumption; *p-value<0.05 is statistically significant; **p-value<0.01 is statistically moderately significant; ***p-value<0.001 is statistically highly significant

Parameters	Group III		Group IV		t-values	p-values
	Mean±SD	SE	Mean±SD	SE		
Age (years)	14.93±0.59	0.153	15.27±0.59	0.15	-1.54	0.135
Height (cm)	168.5±4.06	1.676	167.6±4.20	1.26	2.086	0.081
Weight (Kg)	65±4.94	1.362	68.25±9.75	1.72	2.017	0.091
BMI (kg/m ²)	23.33±1.16	0.363	24.06±2.8	0.44	2.129	0.054*
BSA (m ²)	1.77±0.1	0.027	1.74±0.7	0.07	1.867	0.128
PR (bpm)	73.58±8.37	3.798	74.86±9.86	3.09	2.979	0.025*
SBP (mmHg)	121.1±14.52	3.749	127.1±11.8	3.06	-1.240	0.225
DBP (mmHg)	68.93±8.24	2.128	72.13±7.2	1.85	-1.135	0.266
PP (mmHg)	48.5±5.31	3.721	51.2±6.3	2.75	4.027	0.02*
MAP (mmHg)	87.6±6.81	2.151	87.9±5.13	1.66	3.488	0.021*
WC (cm)	71.47±5.82	1.502	82.53±3.87	1	-6.13	<0.001
HC (cm)	90.13±3.72	0.96	97.1±7.23	1.87	-3.32	0.003**
WHR	0.79±0.04	0.011	0.85±0.08	0.02	-2.68	0.012*
RR (cpm)	18.27±2.49	0.64	19.13±2.29	0.59	-0.99	0.33
BF%	21.55±2.23	0.575	28.7±2.63	0.68	-8.02	<0.001
FM (kg)	10.44±2.05	0.528	16.72±3.28	0.85	-6.29	<0.001
FFM (kg)	37.66±3.47	0.897	41.11±3.7	0.9	-2.64	0.014*
MM (%)	18.83±1.74	0.449	20.6±1.85	0.48	-2.64	0.014*
PFI	52.06±4.83	1.248	45.84±3.39	0.88	4.079	<0.001
VO ₂ max (mL/kg/min)	59.64±3.53	0.912	53.33±3.33	0.86	5.036	<0.001


[Table/Fig-2]: Unpaired t-test results of anthropometric and physiological characteristics between Group III (normal weight girls) and Group IV (overweight/obese girls). Data are Mean±SD and SE; Values in the final column represent results of unpaired t-test among two groups; Significant difference between groups showed at p<0.05 level; BMI: Body mass index; BSA: Body surface area; SBP: Systolic blood pressure; DBP: Diastolic blood pressure, PP: pulse pressure, MAP: Mean arterial pressure; WC: Waist circumference; HC: Hip circumference; PR: Pulse rate; SBP: systolic blood pressure; DBP: Diastolic blood pressure, PP: Pulse pressure; MAP: Mean arterial pressure; WHR: Waist hip ratio; BF %: Body fat percentage, FM: Fat mass; FFM: Fat-free mass; FMI: Fat mass index; MM: Muscle mass; PFI: Physical fitness index and VO₂max: Maximal oxygen consumption; *p-value<0.05 is statistically significant; ** p-value<0.01 is statistically moderately significant; *** p-value<0.001 is statistically highly significant

[Table/Fig-3,4] depict high values of PFI and VO₂max among Group I (normal weight boys) compared to Group II (overweight/obese boys) Group III (normal weight girls) compared to Group IV (overweight/obese girls).


[Table/Fig-5,6] showed Pearson correlation between BMI and PFI also BMI and VO₂max. PFI (p-value<0.001; r=-0.504) and VO₂max (p-value<0.001; r=-0.459) showed significant negative correlation with BMI.




[Table/Fig-3]: Comparison of cardiopulmonary fitness Parameters between Groups I (Non-obese boys) and II (obese boys).



[Table/Fig-4]: Comparison of cardiopulmonary fitness parameters between groups III (Non-obese girls) and IV (obese girls).



[Table/Fig-5]: Pearson's correlation between physical fitness index (PFI in %) and Body Mass Index (BMI in kg/m²) in both boys and girls.



[Table/Fig-6]: Pearson's correlation between physical fitness index (VO2max) and Body Mass Index (BMI in kg/m²) in both boys and girls.

DISCUSSION

Creating a lifestyle pattern of regular physical activity that will carry forward to adulthood is the best primary prevention

strategy. The clinical management of obese children has proved to be complicated. Present results showed significant negative correlation between PFI (p<0.001; r=-0.504) and VO₂max (p<0.001; r=-0.459) showed with BMI. Our results related to PFI and VO₂max with BMI corroborate with Ortega FB et al., and Patkar KU et al., who reported a higher BMI in adolescents with lower cardiopulmonary fitness independent of their passive and leisure time activities (physical activities outside school) [30,31].

A study showed about 10% of the variance in WC in boys and 18% of the WC variance in girls explained by sedentary activities like television viewing and video/computer also another study showed cardiopulmonary fitness was inversely related to WC (boys: p-value=0.001; girls: p-value=0.005) independent of physical activity. Ortega FB et al., and Patkar KU and Joshi AS showed an inverse relationship between physical fitness and overweight and overweight and physical fitness, except Huotari P et al., where no data was available [30-32]. Excess weight typically raises blood pressure, and weight loss usually lowers blood pressure. Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association [33].

Most of the anthropometric and body composition parameters were higher in the overweight/obese when compared to the standard weight group because of more significant fat content. Moreover, our observation that PFI and VO₂max are inversely related to BMI correlates with multiple studies that have been conducted earlier [30,31]. A survey by Eddolls WTB et al., though structural equation modelling method postulated that enhancing cardiorespiratory fitness and BMI through increasing vigorous physical activity may be of benefit in direct and indirect ways to both Quality Of Life (QoL) and mental well-being in adolescents [34]. Dhanakshirur GB et al., showed that regular exercise and nutrition among school children increase physical fitness compared to sedentary lifestyles [35]. So, authors can suggest: 1) School aged children and youth must accumulate at least 60 minutes of Moderate to Vigorous intensity Physical Activity (MVPA) daily for health benefits, according to the WHO recommendations [36]; 2) The school setting is the most influential, ideal environment for physical activity interventions on children during their early years; 3) Advantages of encouraging physical activity in schools are excelling in academics, lower drop-out rates, and better behavior in class, improved self-confidence and active participation in curricular and extra co-curricular activities; 4) Interventions suggested: incorporation into the school curricula of the correlation between physical activity and nutrition; 5) Well-constructed and implemented the physical education curriculum. Training of staff must accordingly be undertaken; 6) Provision of healthy food in the school cafeteria; 7) Parental involvement could be an integral part of school-based interventions [37]. The number of deaths related to sedentary living or obesity is approximately a half-million per year. Regular physical activity may influence the QoL [38].

Limitation(s)

The study sample size was small.

CONCLUSION(S)

The present study showed that exercise testing might help to identify obese children who are not readily apparent in reduced activity or more cardiac or pulmonary impairment. Thus exercise testing can alert the clinician to children whose obesity may lead to severe cardiopulmonary morbidity and require more vigorous intervention. This study may serve as an eye opener for schools to stress regular physical activity through sports and physical training programs for school children throughout the country to counteract a sedentary lifestyle. Improvements in cardiopulmonary fitness have positive effects on depression, anxiety, mood status, and self

esteem and are associated with higher academic performance. Preventive and social medicine staff should work closely with the policymakers to increase resources for the promotion of physical activity within the school system.

Acknowledgement

Authors would like to thank BLDE (DU) for this opportunity to conduct the research and financial support.

REFERENCES

- [1] Woodard MT. Ageless Mirrorathlete: Overweight and Unfit No More. iUniverse; 2019 Jan 21.
- [2] Marshall SJ. Developing countries face double burden of disease. Bulletin of the World Health Organization. 2004;82:556.
- [3] Blair SN, Kohl HW, Paffenbarger RS, Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality: a prospective study of healthy men and women. JAMA. 1989;262(17):2395-401.
- [4] Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. New England Journal of Medicine. 2002;346(11):793-801.
- [5] Mora S, Redberg RF, Cui Y, Whiteman MK, Flaws JA, Sharrett AR, et al. Ability of exercise testing to predict cardiovascular and all-cause death in asymptomatic women: a 20-year follow-up of the lipid research clinics prevalence study. JAMA. 2003;290(12):1600-07.
- [6] Obesity and overweight. https://www.who.int/health-topics/obesity#tab=tab_1
- [7] Pi-Sunyer X. The medical risks of obesity. Postgraduate Medicine. 2009;121(6):21-33.
- [8] Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. Medicine and Science in Sports and Exercise. 2000;32(5):963-75.
- [9] Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. Circulation. 2007;116(9):1081.
- [10] Trost SG, Pate RR, Saunders R, Ward DS, Dowda M, Felton G. A prospective study of the determinants of physical activity in rural fifth-grade children. Preventive Medicine. 1997;26(2):257-63.
- [11] Pate RR, Trost SG, Felton GM, Ward DS, Dowda M, Saunders R. Correlates of physical activity behavior in rural youth. Research Quarterly for Exercise and Sport. 1997;68(3):241-48.
- [12] Dobbins M, Husson H, DeCorby K, LaRocca RL. School-based physical activity programs for promoting physical activity and fitness in children and adolescents aged 6 to 18. Cochrane Database of Systematic Reviews. 2013;(2).
- [13] Wang LY, Gutin B, Barbeau P, Moore J, Hanes J, Johnson M, et al. Cost-effectiveness of a school-based obesity prevention program. Journal of School Health. 2008;78(12):619-24.
- [14] Taylor HL, Buskirk ER, Henschel A. Maximal oxygen intake as an objective measure of cardio-respiratory performance. J Appl Physiol. 1955;8(1):73-80.
- [15] Lauersen JB, Brønd JC, Anderssen SA, Sardinha LB, Steene-Johannessen J, McMurray RG, et al. A new approach to define and diagnose cardiometabolic disorder in children. J Diabetes Res. 2015;2015:539835.
- [16] Eisenmann JC, Welk GJ, Ihmels M, Dollman J. Fatness, fitness, and cardiovascular disease risk factors in children and adolescents. Med Sci Sports Exerc. 2007;39(8):1251-56.
- [17] Sheldon MR. Introduction to probability and statistics for engineers and scientists. Wiley publication, Newyork, Burkleynedn 1987, Pp.106.
- [18] Ortega RM, Requejo AM, Lopez-Sobaler AM, Quintas ME, Andres P, Redondo MR, et al. Difference in the breakfast habits of overweight/obese and normal weight schoolchildren. International journal for vitamin and nutrition research. Internationale Zeitschrift für Vitamin-und Ernährungsforschung. Journal international de vitaminologie et de nutrition. 1998;68(2):125-32.
- [19] Jain AK. Manual of Practical Physiology. 2nd ed. Himachal Pradesh: Arya Publications; 2007: Pp.178-184,223,233,148-151.
- [20] Pal GK. Text book of Practical Physiology. Ed. Chennai: Orient Longmann Publications; 2005: Pp. 178,154 -160.
- [21] Choudhari AR. Text book of practical physiology. 1st ed. India: Paras Publishers; 2000: Pp. 200-07.
- [22] Wanger J. Pulmonary function testing A practical approach. 1st ed. Baltimore: Williams & Wilkins; 1992.
- [23] Keys A, Fidanza F, Karvonen MJ, Kimura N, Taylor HL. Indices of relative weight and obesity. J Chronic Dis. 1972;25(6):329-43.
- [24] Seidel HM, Ball JW, Dains JE, Flynn JA, Solomon BS, Stewart RW. Mosby's Guide to Physical Examination. 7th ed. St. Louis, MO, USA: Mosby; 2011.
- [25] Deurenberg P, Weststrate JA, Seidell JC. Body mass index as a measure of body fatness: age- and sex-specific prediction formulas. Br J Nutr. 1991;65(2):105-14.
- [26] Cotes JE, Chinn DJ. Body mass, fat percentage, and fat free mass as reference variables for lung function: effects on terms for age and sex. Thorax. 2001;56:839-44.
- [27] Meisner JS, Lawrie JA. Human biology – A guide to field methods. 2nd ed. Oxford: Blackwell Publishers; 1969: Pp.325-328.
- [28] McArdle WD. Energy transfer. In: Essential of exercise Physiology. 3rd ed. Baltimore: Lippincott Williams & Wilkins; 2000: Pp.245.
- [29] Vivian H, Heyward. Table Reference for the Physical Fitness, Specialist certification Manual. 3rd ed. The Cooper Institute for Aerobics Research; 1998: 48.
- [30] Ortega FB, Tresaco B, Ruiz JR, Moreno LA, Martin-Matillas M, Mesa JL, et al. Cardiorespiratory fitness and sedentary activities are associated with adiposity in adolescents. Obesity (Silver Spring, Md). 2007;15(6):1589-99.
- [31] Patkar KU, Joshi AS. Comparison of VO2max in obese and non-obese young Indian population. Indian Journal of Physiology and Pharmacology. 2011;55(2):188-92.
- [32] Huotari P, Nupponen H, Mikkelsen L, Laakso L, Kujala U. Adolescent physical fitness and activity as predictors of adulthood activity. Journal of Sports Sciences. 2011;29(11):1135-41.
- [33] Appel LJ, Brands MW, Daniels SR, Karanja N, Elmer PJ, Sacks FM, American Heart Association. Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association. Hypertension. 2006;47(2):296-308.
- [34] Eddolls WTB, McNarry MA, Lester L, Winn CON, Stratton G, Mackintosh KA. The association between physical activity, fitness and body mass index on mental well-being and quality of life in adolescents. Qual Life Res. 2018;27(9):2313-20.
- [35] Dhanakshirur GB, Aithala M, Khodnapur JP. Role of exercise and nutrition on cardiopulmonary fitness. Indian Medical Gazette. 2012;145(1):06-09.
- [36] WHO. Global Recommendations on Physical Activity for Health. Geneva, Switzerland: World Health Organization; 2010.
- [37] Salas C, Petermann-Rocha F, Celis-Morales C, Martínez-López EJ. Parental support for physical activity in schoolchildren and its influence on nutritional status and fitness. Rev Chil Pediatr. 2018;89(6):732-40.
- [38] Staurowsky EJ, De Souza MJ, Miller KE, Sabo D, Shakib S, Theberge N, et al. Her life depends on it III: sport, physical activity, and the health and well-being of American girls and women. Women's Sports Foundation. 2015 May.

PARTICULARS OF CONTRIBUTORS:

1. Student, Department of Physiology, BLDE (Deemed to be University) Shri B. M. Patil Medical College, Hospital and Research Centre, Vijayapura, Karnataka, India.
2. Associate Professor, Department of Physiology, BLDE (Deemed to be University) Shri B. M. Patil Medical College, Hospital and Research Centre, Vijayapura, Karnataka, India.
3. Assistant Professor, Department of Physiology, SDM College of Medical Sciences and Hospital, Dharawad, Karnataka, India.
4. Professor and Head, Department of Physiology, BLDE (Deemed to be University) Shri B. M. Patil Medical College, Hospital and Research Centre, Vijayapura, Karnataka, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Jyoti P Khodnapur,
Associate Professor, Department of Physiology, BLDE (Deemed to be University),
Shri B. M. Patil Medical College, Hospital and Research Centre,
Vijayapura-586103, Karnataka, India.
E-mail: jyoti.khodnapur@bldedu.ac.in

PLAGIARISM CHECKING METHODS: [Jan H et al.]

- Plagiarism X-checker: Mar 12, 2021
- Manual Googling: Jun 30, 2021
- iThenticate Software: Aug 14, 2021 (25%)

ETYMOLOGY: Author Origin

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. NA

Date of Submission: **Mar 06, 2021**

Date of Peer Review: **May 08, 2021**

Date of Acceptance: **Jul 15, 2021**

Date of Publishing: **Oct 01, 2021**