

**“SERUM URIC ACID LEVELS AND SERUM LIPID  
LEVELS IN PATIENTS WITH ISCHAEMIC  
CEREBROVASCULAR ACCIDENT”**

**By**

**Dr. TAVISH ARORA**

Dissertation submitted to

**BLDE University, Vijayapur, Karnataka**



In partial fulfilment of the requirements for the degree of

**MD**

**IN**

**GENERAL MEDICINE**

Under the guidance of

**Dr. P. G. MANTUR** M.D

Associate Professor

**DEPARTMENT OF MEDICINE,**

BLDEU'S Shri B.M. Patil Medical College,

Vijayapur, Karnataka

**2017**

# BLDE UNIVERSITY, VIJAYAPUR, KARNATAKA



## DECLARATION BY THE CANDIDATE

I hereby declare that this dissertation/thesis entitled “**SERUM URIC ACID LEVELS AND SERUM LIPID LEVELS IN PATIENTS WITH ISCHAEMIC CEREBROVASCULAR ACCIDENT** ” is a bonafide and genuine research work carried out by me under the guidance of **DR. P.G. MANTUR<sub>M.D.</sub>**, Associate Professor, Department of Medicine, Shri B.M. Patil Medical College, Vijayapur.

Date:

Place: Vijayapur

**Dr. TAVISH ARORA**

**BLDE UNIVERSITY, VIJAYAPUR, KARNATAKA**



**CERTIFICATE BY THE GUIDE**

This is to certify that the dissertation entitled “**SERUM URIC ACID LEVELS AND SERUM LIPID LEVELS IN PATIENTS WITH ISCHAEMIC CEREBROVASCULAR ACCIDENT** ” is a bonafide and genuine research work carried out by **Dr. TAVISH ARORA** in partial fulfilment of the requirement for the degree of MD in General medicine.

Date:

Place: Vijayapur

**Dr. P.G. MANTUR** M.D.  
Associate Professor  
Department of Medicine,  
Shri B.M. Patil Medical College,  
Vijayapur.

**BLDE UNIVERSITY, VIJAYAPUR, KARNATAKA**



**ENDORSEMENT BY THE HOD, PRINCIPAL / HEAD OF THE  
INSTITUTION**

This is to certify that the dissertation entitled “**SERUM URIC ACID LEVELS AND SERUM LIPID LEVELS IN PATIENTS WITH ISCHAEMIC CEREBROVASCULAR ACCIDENT**” is a bonafide research work done by **Dr. TAVISH ARORA** under the guidance of, **Dr. P.G. MANTUR** M.D., Associate Professor, Department of Medicine, Shri. B.M. Patil Medical College, Vijayapur.

Seal & Signature of  
HOD of Medicine  
**Dr. M.S. MULIMANI** MD

BLDEU’s Shri B.M. Patil  
Medical College, Hospital &  
Research Centre, Vijayapur.

Date:

Place: Vijayapur

Seal and signature of  
the principal  
**DR. S.P. GUGGARIGUDAR** MS

BLDEU’s Shri B.M. Patil  
Medical College, Hospital &  
Research Centre, Vijayapur.

Date:

Place: Vijayapur

# BLDE UNIVERSITY, VIJAYAPUR, KARNATAKA



## COPYRIGHT

### DECLARATION BY THE CANDIDATE

I hereby declare that the BLDE University, Karnataka shall have the rights to preserve, use and disseminate this dissertation / thesis in print or electronic format for academic / research purpose.

Date:

Place: Vijayapur

**Dr. TAVISH ARORA**

© BLDE University, Vijayapur, Karnataka

## ACKNOWLEDGEMENT

I wish to express my deep sense of gratitude and regards to my guide **Dr. P.G. Mantur MD**, Associate Professor of Medicine, for his able guidance and valuable suggestions, constant supervision and encouragement which he rendered in pursuit of my postgraduate studies and during the preparation of this dissertation.

I would also like to express my sincere thanks to our vice-chancellor and Professor **Dr. M. S. Biradar MD**, for his kind support.

I would also like to express my sincere thanks to our principal **Dr. S. P. Guggarigoudar MS**, for his kind support.

I am extremely grateful to Professors of Medicine, **Dr. R. C. Bidri MD Dr. S.N. Bentoor MD, Dr. S. S. Devarmani MD, Dr. L. S. Patil MD, Dr. S.R. Badiger MD & Dr. M. S. Mulimani MD** for their valuable advice and guidance in completing this work.

I thank **Dr. Shahnawaz**, statistician who kindly obliged & helped me with the statistical work and I thank all my friends and colleagues for their kind cooperation.

I would also like to thank my parents **Mr. Surendra Singh Arora** and **Mrs. Parveen Arora**, without their constant encouragement and moral support, my studies would have been a distant dream.

I would like to thank the Almighty god who gave me the energy, skill and the enthusiasm to complete this as well as the other tasks in my life & also for continuing to shower his blessings upon me.

Last but not the least; I extend my sincere thanks to all the patients who willingly cooperated in this study.

**Date:**

**Dr. TAVISH ARORA**

**Place: Vijayapur**

## **LIST OF ABBREVIATIONS**

|      |   |                              |
|------|---|------------------------------|
| TC   | - | Total cholesterol            |
| VLDL | - | Very low density lipoprotein |
| LDL  | - | Low density lipoprotein      |
| TG   | - | Triglycerides                |
| HDL  | - | High density lipoprotein     |
| CAD  | - | Coronary artery disease      |
| SUA  | - | Serum uric acid              |
| CBC  | - | Complete blood count         |
| CT   | - | Computerised tomography      |
| MRI  | - | Magnetic resonance imaging   |
| RBS  | - | Random blood sugar           |
| ECG  | - | Electrocardiogram            |

## **ABSTRACT**

### **BACKGROUND**

Stroke is a growing disease and it is the second common cause of death in the world after coronary heart disease especially in the elderly. Uric acid is the ultimate catabolite of purine metabolism in human and higher primates. It exists in the extracellular compartment as sodium urate, and it is cleared from the plasma through the kidney. In patients with acute stroke hyperuricemia was significantly higher than normal population. Hyperuricemia has been associated with increased amounts of triglycerides and LDL cholesterol.

### **OBJECTIVES:**

Serum uric acid and lipid levels in patients with ischaemic cerebrovascular accident.

### **METHODOLOGY:**

The parameters were collected from the patients attending the outpatient and inpatient departments in BLDEU'S Shri. B.M. Patil Medical College Hospital and Research Centre, Vijayapur from January 2015 to June 2016.

### **RESULTS:**

A total of 60 patients with ischemic stroke were included in the study. Out of which 43 (71.7%) were males and 17 (28.3%) were females. The mean age of the patients was  $63.2 \pm 14.8$ . Mean serum uric acid levels in the patients studied was  $5.5 \pm 1.7$ , and 18 patients (30%) were hyperuricemic. Serum uric acid levels were significantly higher in females ( $6.2 \pm 1.7$ ) compared to the males ( $5.2 \pm 1.6$ ). Dyslipidemia was seen in 49 (81.7%) patients (Males were 36 and females were 13). It was predominant in the age group between 56-70 years. More than one lipid

parameter was seen to be deranged in 10 male patients and 4 female patients. Of them 87.75% of patients had low HDL levels, 36.73%, 10.20% and 8.16% patient had high cholesterol, triglycerides and LDL levels respectively. The study also showed that 15 patients had both hyperuricemia and dyslipidemia.

## **CONCLUSION**

Hyperuricemia and its accompanying dyslipidemia can be considered as the risk factor for acute ischaemic stroke.

## **KEYWORDS:**

Uric acid, Acute stroke, HDL, LDL, TC, TG.

## TABLE OF CONTENTS

| <b>Sl. No</b> | <b>Contents</b>             | <b>Page No</b> |
|---------------|-----------------------------|----------------|
| <b>1</b>      | <b>INTRODUCTION</b>         | <b>1</b>       |
| <b>2</b>      | <b>AIMS AND OBJECTIVES</b>  | <b>2</b>       |
| <b>3</b>      | <b>REVIEW OF LITERATURE</b> | <b>3</b>       |
| <b>4</b>      | <b>METHODOLOGY</b>          | <b>32</b>      |
| <b>5</b>      | <b>RESULTS</b>              | <b>34</b>      |
| <b>6</b>      | <b>DISCUSSION</b>           | <b>49</b>      |
| <b>7</b>      | <b>CONCLUSION</b>           | <b>51</b>      |
| <b>8</b>      | <b>SUMMARY</b>              | <b>53</b>      |
| <b>9</b>      | <b>BIBLIOGRAPHY</b>         | <b>54</b>      |
| <b>10</b>     | <b>ANNEXURES</b>            |                |
|               | ETHICAL CLEARANCE           | 61             |
|               | INFORMED CONSENT FORM       | 62             |
|               | PROFORMA                    | 63             |
|               | KEY TO MASTER CHART         | 67             |
|               | MASTER CHART                | 68             |

## LIST OF TABLES

| Sl. No. | Tables  | Page No. |
|---------|---|----------|
| 1       | Distribution of cases according to Sex                      | 34       |
| 2       | Distribution of cases according to Age (Yrs)                | 34       |
| 3       | Mean age by Sex   | 35       |
| 4       | Distribution of cases according to Deficit                  | 36       |
| 5       | Distribution of cases according to Hyperuricemia            | 37       |
| 6       | Mean Distribution of Serum Uric Acid according to Sex       | 38       |
| 7       | Hyperuricemia and Sex                                       | 39       |
| 8       | Mean Distribution of Serum Uric Acid according to Age (Yrs) | 40       |
| 9       | Hyperuricemia and Age (Yrs)                                 | 41       |
| 10      | Correlation of Serum Uric Acid with Selected parameters     | 42       |
| 11      | Distribution of cases according to Lipid profile            | 42       |
| 12      | Dyslipidemia and Age  | 43       |
| 13      | Dyslipidemia and Sex  | 44       |
| 14      | Patient having both Hyperuricemia and Dyslipidemia          | 45       |
| 15      | Abnormal Lipid Profile and Hyperuricemia                    | 46       |
| 16      | Abnormal Lipid Profile and Sex                              | 47       |
| 17      | Abnormal Lipid Profile and Age(Yrs)                         | 48       |

## LIST OF GRAPHS

| Sl. No. | Figures   | Page No. |
|---------|---|----------|
| 1       | Distribution of cases according to Sex                      | 34       |
| 2       | Distribution of cases according to Age (Yrs)                | 35       |
| 3       | Mean age by Sex   | 36       |
| 4       | Distribution of cases according to Deficit                  | 36       |
| 5       | Distribution of cases according to Hyperuricemia            | 37       |
| 6       | Mean Distribution of Serum Uric Acid according to Sex       | 38       |
| 7       | Hyperuricemia and Sex                                       | 39       |
| 8       | Mean Distribution of Serum Uric Acid according to Age (Yrs) | 40       |
| 9       | Hyperuricemia and Age (Yrs)                                 | 41       |
| 10      | Distribution of cases according to Lipid profile            | 42       |
| 11      | Dyslipidemia and Age  | 43       |
| 12      | Dyslipidemia and Sex  | 44       |
| 13      | Patient having both Hyperuricemia and Dyslipidemia          | 45       |
| 14      | Abnormal Lipid Profile and Hyperuricemia                    | 46       |
| 15      | Abnormal Lipid Profile and Sex                              | 47       |
| 16      | Abnormal Lipid Profile and Age(Yrs)                         | 48       |

# INTRODUCTION

A stroke or cerebrovascular accident is defined as an abrupt onset of a neurological deficit that is attributable to a focal vascular cause. A definition of stroke is clinical and laboratory studies including brain imaging are used to support the diagnosis. The clinical manifestations of stroke are highly variable because of the complex anatomy of the brain and its vasculature. Cerebral ischemia is caused by reduction in blood flow that last longer than several seconds.<sup>1</sup>

Stroke is the second leading cause of death worldwide and it is also one of the leading causes of adult disability.<sup>2</sup> Numerous risk factors are involved in the development of stroke such as hypertension, smoking, dyslipidemia and diabetes mellitus. Hyperuricemia has been reported to be an independent predictor of stroke.<sup>3</sup>

There is a pressing need to identify these treatable risk factors that can be easily measured and are highly prevalent, in order to identify patients at high risk for stroke.

Hyperuricemia have also been suggested as one of the factors in the pathogenesis of an atheroma. Significant association was found between serum uric acid and serum triglycerides. This implicates that a rise in serum uric acid and serum triglyceride may play some part in the etiology of ischemic cerebrovascular disease.<sup>4</sup>

But contrary to this, other studies have also advocated uric acid to be neuroprotective due to its anti-oxidation action. Considering these conflicting data, we undertook this study to evaluate serum uric acid and serum lipid levels in patients with ischaemic cerebrovascular stroke.

## **AIMS AND OBJECTIVES**

Serum uric acid levels and serum lipid levels in patients with ischaemic cerebrovascular accident.

## REVIEW OF LITERATURE

Uric acid is the ultimate catabolite of purine metabolism in human and higher primates. It exists in the extracellular compartment as sodium urate, and it is cleared from the plasma through the kidney. It has been reported that increased levels of uric acid are associated with established cardiovascular risk factor such as elevated serum triglyceride and cholesterol concentration, hypertension, obesity, insulin resistance and metabolic syndrome.<sup>5-9</sup>

Uric acid is the breakdown product of purines. In this process hypoxanthine is converted by the enzyme xanthine oxidase to xanthine and further to uric acid. Both steps induce the release of free radicals. Increased uric acid levels promote oxygenation of low-density lipoprotein cholesterol and facilitate lipid peroxidation. Uric acid may stimulate vascular smooth cell proliferation, and reduce vascular nitric oxide production. Moreover, higher uric acid levels may be associated with increased platelet adhesiveness predisposing to thrombus formation. SUA has also been found to stimulate the synthesis of pro-inflammatory factors like monocyte chemoattractant protein-1, interleukin-1 $\beta$ , interleukin-6, and tumor necrosis factor- $\alpha$ . Experimental findings indicate that uric acid might have a role in the development of hypertension through stimulation of the renin-angiotensin system and induction of sodium sensitivity. In rats, uric acid has been shown to mediate renal disease development by causing glomerular hypertension and hence renal hypertrophy, glomerulosclerosis, and interstitial fibrosis. Uric acid also induces renal arteriolar thickening independently of its effect on blood pressure. Each of these factors can play a pivotal role in the progression of atherosclerosis. As compared to control coronary artery walls, urate crystals are more abundant in diseased atherosclerotic plaques. It has been

suggested that serum uric acid may cause endothelial dysfunction. Even a mild elevation of serum uric acid was associated with cerebral ischemia in adults. It was suggested that impaired vascular tone and endothelial dysfunction could contribute to ischemic changes, because they permit cerebrospinal fluid to cross the blood-brain barrier and cause areas of edema.<sup>10-21</sup>

## **LIPIDS AND ATHEROSCLEROSIS**

Several evidences have contributed to our current understanding of the relationship between increase in plasma cholesterol and development of CAD. Premature atherosclerosis results from high cholesterol levels, even in the absence of other cardiovascular risk factors.

The onset of atherosclerosis occurs early in life with diffuse regular thickening of the arterial intima in childhood. The smooth appearance of the arterial tree is usually lost during the teenage years with formation of nodular aggregates or cushions of fibro-elastic tissue, termed fatty streaks.<sup>22</sup>


Fatty streaks are collections of lipid, mainly cholesterol esters in macrophages and smooth muscle cells deposited in the intima of the artery. These fatty streaks are the precursors of the hallmark of atherosclerosis, the fibrous atheromatous plaque. The fibrous plaques are white lesions that usually protrude into the vessel lumen and consists of a core of cholesterol, cholesterol ester, phospholipid and necrotic cells covered by a fibrous cap of elastin and collagen. There is also proliferation of the smooth muscle cells into the media. Another important component is the foam cells, which contain a large quantity of lipid.<sup>23</sup>

Hypertriglyceridemia also maybe a marker of an individual with a genetic defect of lipoprotein metabolism, such as accumulation of VLDL remnants which are associated with premature atherosclerosis.

An important factor for the development of atheromatous plaques is oxidized LDL, which causes increased chemotaxis of monocytes, and increased uptake of oxidized LDL by macrophages.<sup>24</sup>

## **URIC ACID**

Uric acid is a heterocyclic compound of carbon, nitrogen, oxygen and hydrogen with the formula  $C_5H_4N_4O_3$ . It is the final breakdown product of purine metabolism in humans.



**Figure-1**

## **HISTORY**

In 1776, the Swedish chemist Scheele isolated uric acid from a calculus of urinary tract and in 1797, the British chemist Wollaston isolated the substance from a tophus which he removed from his own ear. 50 years later, these observations led the British Physician, Alred Barry Garrod to show by chemical isolation, the presence of higher concentrations of uric acid in the blood of gouty patients. Garrod's subsequent studies formulated, for the first time, a rational relationship between hyperuricemia and the clinical symptomatology of gouty arthritis. Later, in 1913, Folin F. Denin developed a more reliable method of chemical determination of uric acid which led to rejection of Garrod's concept.<sup>25-29</sup>

## STRUCTURE AND CHEMISTRY




Figure - 2

The chemical nature of uric acid was shown by Fisher to be 2,6,8 trihydroxypurine. Uric acid is a diprotic acid with  $pK_{a1}=5.4$  and  $pK_{a2}=10.3$ . Thus in strong alkali at high pH or in the presence of carbonic acid or carbonate ions, it forms the singly charged hydrogen or acid uric acid ion as its  $pK_{a2}$  is greater than the  $pK_{a1}$  of carbonic acid. As its second ionization is so weak, the full uric acid salts tend to hydrolyze back to hydrogen uric acid salts and free base at pH values around neutral. It is aromatic because of the purine functional group.<sup>30</sup>

## SOLUBILITY

Generally, the water solubility of uric acid and its alkali metal and alkaline earth salts is rather low. All these salts exhibit greater solubility in hot water than cold, allowing for easy recrystallization. This low solubility is significant for the etiology of gout. The solubility of the acid and its salts in ethanol is very low or negligible. In ethanol water mixtures, the solubilities are somewhere between the end values for pure ethanol and pure water.

## BIOLOGY

The enzyme xanthine oxidase makes uric acid from xanthine and hypoxanthine, which in turn are produced from purines. Uric acid is released in hypoxic conditions.<sup>31</sup>

In humans and higher primates, uric acid is the final oxidation (breakdown) product of purine metabolism and is excreted in urine. In most other mammals, the enzyme uricase further oxidizes uric acid to allantoin. The loss of uricase in higher primates parallels the similar loss of the ability to synthesize ascorbic acid, leading to the suggestion that uric acid may partially substitute for ascorbate in such species. Both uric acid and ascorbic acid are strong reducing agents (electron donors) and potent antioxidants.<sup>32</sup>


In humans, over half the antioxidant capacity of blood plasma comes from uric acid. The Dalmatian dog has a genetic defect in uric acid uptake by the liver and kidneys, resulting in decreased conversion to allantoin, so this breed excrete uric acid, and not allantoin, in the urine. In birds and reptiles, and in some desert dwelling mammals (e.g., the kangaroo rat), uric acid also is the end product of purine metabolism, but it is excreted in feces as a dry mass. This involves a complex metabolic pathway that is energetically costly in comparison to processing of other nitrogenous wastes such as urea (from urea cycle) or ammonia, but has the advantage of reducing water loss.

In humans, about 70% of daily uric acid disposal occurs via the kidneys, and in 5-25% of humans, impaired renal (kidney) excretion leads to Hyperuricemia.

## **GENETICS**

A proportion of people have mutations in the proteins responsible for the excretion of uric acid by the kidneys. Nine genes have so far been identified: SLC2A9; ABCG2; SLC17A1; SLC22A11; SLC22A12; SLC16A9; GCKR; LRRC16A; and PDZK1. SLC2A9 is known to transport both uric acid and fructose.<sup>33</sup>

## SYNTHESIS



**Synthesis of Uric Acid Figure-3**

Degradation of Purines and Pyrimidines Produces Uric Acid and Urea, Respectively. Purine nucleotides are degraded by a pathway in which they lose their phosphate through the action of **5'-nucleotidase**. Adenylate yields adenosine, which is deaminated to inosine by **adenosine deaminase**, and inosine is hydrolyzed to hypoxanthine (its purine base) and D-ribose. Hypoxanthine is oxidized successively to xanthine and then uric acid by **xanthine oxidase**, a flavoenzyme with an atom of molybdenum and four iron-sulfur centers in its prosthetic group.

Molecular oxygen is the electron acceptor in this complex reaction. GMP catabolism also yields uric acid as end product. GMP is first hydrolyzed to guanosine, which is then cleaved to free guanine. Guanine undergoes hydrolytic removal of its amino group to yield xanthine, which is converted to uric acid by xanthine oxidase.

### **DIETARY INTAKE**

Diet provides an important exogenous source of purines. Dietary intake of purines, nucleic acids contribute to the serum uric acid level and the daily excretion of uric acid, this contribution being proportional to the intake. However, the magnitude of this component of daily uric acid production is usually relatively small, since serum uric acid levels decrease by only 1 mg/dl or less when an individual changes from a normal to purine free diet. On the other hand, diets unusually high in purines (such as liver, meat sweet breads, kidney and anchovy) have the potential to raise the plasma uric acid concentration significantly. Because about 50% of the ingested RNA purine and 25% of the ingested DNA purine appear in urine as uric acid, foods high in nucleic acid content have a significant effect on the serum uric acid level.

### **NORMAL VALUES**

Usually, the uric acid pool size of an adult male, is about, 1200mg, and 700mg uric acid is produced daily. The production is balanced by the excretion of 500mg of uric acid into the urine and 200mg into the small intestine, Any imbalance in the two, results in hyperuricemia or hypouricemia.

The normal serum values vary with age and sex. Most children have serum uric acid concentrations of 3 to 4 mg/dl. Levels start rising during puberty in males, but remain low in females till menopause. The gender variation could probably be due to higher excretion in females. Normal values in **females are 2.5-5.6 mg/dl** and **males is 3.1-7mg/dl**.

After menopause, values for women increase to approximate those at men. Among adults, concentrations vary with height, body weight, blood pressure and renal function as well as alcohol intake.

### **URIC ACID EXCRETION**

In a normal person on a normal diet about 700-1000 mg of uric acid is produced daily. If a person takes a diet rich in nucleoproteins (meat, particularly glandular meat, meat extracts, legumes) he/she will excrete little excess of uric acid.

On a purine free diet some uric acid is constantly excreted in urine amounting to about 200-500 mg/day. This fraction is referred to as endogenous uric acid. If it is determined for a given individual, the excess above this figure which he/she excretes on a purine containing diet is termed exogenous.

About 70% of uric acid is excreted by the kidneys. The remainder is excreted via the gastro intestinal tract, where it is degraded by uricolytic bacteria to carbon dioxide and ammonia. In patients with renal insufficiency, this extra renal elimination of uric acid may be a major route of disposal.

The mechanisms controlling intestinal elimination of uric acid are not well understood. The amount secreted into the lumen is probably related to the plasma concentration, so that a higher fraction of daily uric acid production may be excreted via gastro intestinal tract in hyperuricemic individuals than in normal. It is not thought that intestinal elimination of uric acid represents an important regulator of the plasma uric acid concentration.<sup>34</sup>

On the other hand, renal excretion is the major regulator of plasma uric acid, decreases or increases in renal clearance are readily reflected by inverse changes in plasma uric acid concentration.

Steele and Rieselbach proposed the four component model of renal handling of uric acid mechanism. It includes-

1. Glomerular filtration
2. Presecretory tubular reabsorption.
3. Tubular secretion
4. Postsecretory reabsorption

Normally, uric acid is totally filtered in the renal glomeruli and then the filtered load is completely reabsorbed in the proximal tubules. 5-10% is later secreted into the distal tubules and excreted in the urine. The filtered uric acid undergoes extensive absorption with at least 98% of the filtered load reabsorbed. Recent evidences suggest that much of the secreted uric acid undergoes extensive reabsorption. All four components are influenced by other factors. Since uric acid binding by plasma protein is thought not to be important, the rate of uric acid filtration should vary directly with changes in glomerular filtration rate. Tubular reabsorption is an active process closely related to sodium reabsorption. Expansion of extracellular fluid volume increases the uric acid clearance by inhibiting its tubular reabsorption. Conversely, contraction of extracellular fluid volume, decreases the uric acid clearance, enhancing its tubular reabsorption, thereby leading to hyperuricemia.

The tubular secretory component is directly proportional to plasma uric acid concentration. Secretion may be decreased, if the transport pathway is competitively inhibited by other organic acids. In summary, normally kidneys excrete an amount equivalent to 5-10% of the filtered load of uric acid.

Altered renal excretion of uric acid exhibiting hyperuricemia could be due to -

1. Reduced filtration of uric acid
2. Enhanced reabsorption.
3. Decreased secretion.

No unequivocal data establish any one of these mechanisms as the basic defect, and it is likely that all three are operative within the hyperuricemic population.

Patients with renal disease usually have a reduction in glomerular filtration rate leading to decreased filtered load of uric acid and thus hyperuricemia.

Tubular reabsorption of uric acid is an active process that is closely related to sodium reabsorption. Therefore states of volume contraction (with enhanced sodium reabsorption) are accompanied by increased uric acid reabsorption and decreased uric acid clearance. Diuretic induced volume depletion leads to enhanced tubular reabsorption of uric acid as well as decreased uric acid filtration.

In some situations hyperuricemia has been attributed to competitive inhibition of uric acid secretion by excess organic acids thought to be secreted by the same renal tubular mechanism responsible for uric acid secretion. Examples include starvation (ketosis and free fatty acids), alcoholic ketosis, diabetic ketoacidosis, maple syrup urine disease and lactic acidosis of any cause.

### **HYPERURICEMIA**

This may either be due to increased production or decreased excretion of uric acid or from a combination of the 2 processes. It is defined as a serum concentration more than 420  $\mu\text{mol/L}$  is 7mg/dl as based on physicochemical, epidemiologic and disease-related criteria, in epidemiologic studies, hyperuricemia is defined as the mean plus 2 standard deviations of values determined from a randomly selected healthy population. When measured, in unselected individuals, 95% have serum uric

acid concentration  $<420\mu\text{mol/L}$  (7mg/dl). In relation to the risk of the disease, uric acid levels  $420\mu\text{mol/L}$  is associated with the risk of gouty arthritis and escalates in proportion to the degree of elevation. Hyperuricemia is reported in 2-13.2% of ambulatory adults.

**CAUSES:**<sup>35-38</sup>

**1) Uric acid overproduction**

- Primary idiopathic
- HGPRT deficiency
- PRPP synthetase overactivity
- Hemolytic processes
- Lymphoproliferative diseases
- Myeloproliferative diseases
- Polycythemia vera
- Psoriasis
- Paget's disease
- Rhabdomyolysis
- Exercise
- Alcohol
- Obesity
- Purine-rich diet
- Glycogen storage diseases (types V, VII)

## 2) URIC ACID UNDER EXCRETION

- Primary idiopathic
- Renal insufficiency
- Polycystic kidney disease
- Diabetes insipidus
- Hypertension
- Acidosis-lactic acidosis, ketoacidosis
- Berylliosis
- Hypothyroidism
- Hyperparathyroidism
- Toxemia of pregnancy
- Lead intoxication
- Bartter's syndrome
- Down's syndrome
- Drugs
  - Salicylates (>2 g/d)
  - Diuretics
  - Alcohol
  - L- dopa
  - Ethambutol
  - Pyrazinamide
  - Nicotinamide
  - Cyclosporine

### 3) COMBINED MECHANISM

- Glucose-6-phosphate Dehydrogenase deficiency
- Alcohol
- Shock
- Fructose-1-phosphate aldolase deficiency.
- Physical exercise
- Status epilepticus
- Myocardial infarction
- Acute respiratory failure

#### **HYPOURICEMIA:**

It is defined as a serum uric acid concentration less than 180  $\mu\text{mol/L}$ ; i.e. 3mg/dl. It may occur as a consequence of decreased formation of uric acid or increased renal clearance of uric acid. It can also result when the activity of xanthine oxidase is reduced, either because of deficiency of the enzyme, as in xanthinuria, or when there is pharmacologic inhibition of xanthine oxidase by allopurinol.

Renal causes of hypouricemia range from isolated defects of uric acid reabsorption to more complex abnormalities of renal tubular function. The amount of uric acid finally excreted in urine appears to depend on the balance between tubular secretion of uric acid and the so-called post- secretory reabsorption of secreted uric acid. In contrast to hyperuricemia, which has been studied extensively, only recent observations have suggested that hypouricemia may prove an important clue for the existence of underlying disease processes.

30 years ago, Ramsdell and Kelley in a short communication, looked at the clinical significance of hypouricemia in a hospital population, but some of their plasma uric acid determinations were performed using the colorimetric

phosphotungstate method which is not as specific for uric acid as the uricase method which is now preferred.<sup>39,40</sup>

In a study done in UK, the prevalence of hypouricemia in hospital population was studied. Prevalence was reported to be 6.5% among the males and 4.8% among the females. The large group displaying hypouricemia, was intensive care patients, possibly due to a combination of alteration in the renal handling of uric acid and reduced dietary intake of protein. The author reports that in the study, a large group of patients (14%) who manifested hypouricemia, were those with diabetes mellitus, 10% of whom were on insulin and 4% on sulphonylureas.<sup>41</sup>

#### **CAUSES OF HYPOURICEMIA:<sup>42</sup>**

- Total parenteral nutrition
- Cirrhosis
- Neoplasms
- Diabetes mellitus
- Syndrome of inappropriate secretion of ADH
- Fanconi syndrome
- Xanthine oxidase deficiency
- Drugs
  - o Ascorbic acid
  - o Dicoumarol
  - o Diflunisal
  - o Sulfinpyrazone
  - o NSAIDs
  - o Probenecid
  - o Estrogen

## **OXIDATIVE STRESS<sup>43</sup>**

UA is a marker of oxidative stress, and may have a potential therapeutic role as an antioxidant .On the other hand, like other strong reducing substances such as ascorbate, UA can also act as a pro oxidant, particularly at elevated levels. Thus, it is unclear whether elevated levels of uric acid in diseases associated with oxidative stress such as stroke and atherosclerosis are a protective response or a primary cause. Some researchers propose Hyperuricemia-induced oxidative stress is a cause of metabolic syndrome. On the other hand, plasma uric acid levels correlate with longevity. In primates and other mammals. This is presumably a function of uric acid's antioxidant properties.

## **INDICATIONS FOR URIC ACID LOWERING THERAPY:**

- Initiation of chemotherapy or radiation therapy.
- History of kidney stones.
- History of gouty attacks (>2 per year), tophi.
- Moderate Renal function impairment (creatinine clearance <42 mL/min)  
Renal dysfunction.
- Markedly elevated uric acid levels (12 mg/dL in men or women).

Hyperuricemia is a biochemical abnormality, Not a disease. Indications for lowering uric acid is clear. Drug therapy to lower UA (ULT) for CVD prevention is not advised as per current evidence.

## **URIC ACID ESTIMATION <sup>44</sup>.**

The method used for analysis is enzymatic method (Uricase method ) by using auto analyzers. In our laboratory, values taken as normal range is as follows :

Males 3.1 – 7.0 mg/dl

Females 2.5 – 5.6 mg/dl

Methods using Uricase, the enzyme that catalyses the oxidation of uric acid to allantoin are most specific. The simplest of these methods measures the differential absorption of uric acid and allantoin at 293nm. The differential absorption before and after incubation with Uricase is proportional to uric acid concentration. This method has been proposed as candidate reference method, and is the most specific method.

### **BRIEF ACCOUNT OF THE LIPID CHEMISTRY**

The lipids are a heterogeneous group of compounds including fats, oils, steroids, waxes and related compounds that are related more by their physical than chemical properties.

The lipids have following functions.

1. They are important dietary constituents not only because of their high energy value, but also because of the fat soluble vitamins and essential fatty acids contained in the fat of natural foods.
2. Fat is stored in adipose tissue, where it also serves as a thermal insulator in the subcutaneous tissues and around certain organs.
3. Non polar lipids act as electrical insulators, allowing rapid propagation of depolarization waves along myelinated nerves.
4. Combination of lipid and protein (lipoproteins) serve as the means of transporting lipids in the blood.

**The lipids are classified as given below.<sup>45</sup>**

#### **A) Simple lipids** (esters of fatty acid with various alcohols)

- Fats: esters of fatty acids with glycerol. Oils are fats in liquid state
- Waxes: esters of fatty acids with higher molecular weight monohydric alcohols.

**B) Complex lipids:** These are esters of fatty acids containing groups in addition to an alcohol and a fatty acid groups)

- Phospholipids: lipids containing in addition to fatty acids and an alcohol, a phosphoric acid residue.
- Glycolipids: lipids containing a fatty acid, sphingosine and carbohydrate.
- Other complex lipids: lipids such as sulfolipids and aminolipids. Lipoproteins may also be placed in this category.

**C) Derived lipids:** These include fatty acids, glycerol, steroids, other alcohols, fatty aldehydes, hydrocarbons, lipid soluble vitamins and hormones.

## **FATTY ACIDS**

They are present as such in minute concentrations in plasma cells. Fatty acids occur mainly as esters in natural fats and oils but do occur in the unesterified form as free fatty acids, a transport form found in the plasma. Fatty acids that occur in natural fats are usually straight chain derivatives containing an even number of carbon atoms.

### **Lipid Chemistry**

Fatty acids are of two types

- a. Saturated fatty acids - Those which contain no double bonds.
- b. Unsaturated fatty acids- Those which contain one or more double bonds.

Saturated fatty acids having 10 or less carbon atoms are called "Lower fatty acids"

e.g. - Acetic acid, butyric acid

Saturated fatty acids having more than 10 carbon atoms are

Called "higher fatty acids". e.g. - Palmitic acid, stearic acid.

Unsaturated fatty acids are classified according to degree of saturation.

1. **Mono unsaturated fatty acids** - These are those fatty acids which contain one double bond. E.g. Oleic acid.

2. **Polyunsaturated fatty acids** - These are fatty acids which contain more than one double bond.

eg. Linoleic acid series

Linolenic acid series

Arachidonic acid series

Essential fatty acids are those which cannot be synthesized in body and must be provided in the diet. Lack of these essential fatty acids in diet can produce growth retardation and other deficiency syndromes. Free fatty acids are immediately available energy source and provide much of the energy requirements of body. Normal value ranges from 250 - 400 mg / dl.

## **CHOLESTEROL**

Cholesterol is widely distributed in all cells of the body. It is the best known steroid because of its association with atherosclerosis and heart disease.

It occurs as a white or faintly yellow, almost odorless, pearly leaflets or granules. It is insoluble in water. Sparingly soluble in alcohol and soluble in ether, chloroform, hot alcohol, ethyl acetate alcohol and vegetable "oil.

Cholesterol is found in largest amounts in normal human adult brain and nervous tissue of about 20%, in liver - 0.3%, skin - 0.3%, intestinal mucosa 0.2% and certain endocrine glands namely adrenal cortex contain about 10%.

The normal level of serum total cholesterol in adults varies from 150-250 mg/dl

## **TRIGLYCERIDES (NEUTRAL FATS)**

These molecules are used to provide energy. In the body, stored fat in adipose tissue is the storage form of energy.

Important sites of adipose tissue are subcutaneous tissue around some internal organs and omentum. Fat under the skin prevent heat loss in winter and the intestinal organs get support from fat around them. The triglycerides constitute the body's main caloric reserve. Normal value ranges from 40-150 mg %.

## **PHOSPHOLIPIDS**

Phospholipids are compound lipids. They contain in addition to fatty acids and glycerol one more alcohol or phosphatidic acid, residue nitrogen containing base and other substituents.

Phosphatidic acid is important as an intermediate in the synthesis of triacylglycerols as well as phosphoglycerols but is not found in any great quantity in tissues.

They are classified into 3 groups.

1. **Glycerophospholipid** - Here glycerol is the alcohol group.

eg - Lecithin, Cephalin

2. **Phospho inositides** - Here inositol is the alcohol group.

eg - phosphatidylinositol

3. **Sphingophospholipids** - Here sphingol is the alcohol group.

eg - Sphingomyelin

The lecithins are the most abundant phospholipids at the cell membrane and represent a large portion of body's store of choline.

Dipalmitoyl lecithin is a very effective surface active agent and a major constituent of the surfactant preventing adherence, due to surface tension of the inner surface of lungs.

Phosphatidylethanolamine (cephalin) and phosphatidyl serine differ from phosphatidylcholine only in that ethanolamine or serine, respectively replaces choline.

Sphingomyelins are found in large quantities in brain and nerve tissue.

On hydrolysis, the sphingomyelins yield a fatty acid, phosphoric acid, choline and a complex amino alcohol, sphingosine. No glycerol is present. The combination of sphingosine plus fatty acid is known as ceramide, a structure also found in the glycopospholipids.

### **GLYCOLIPIDS**

Glycolipids are widely distributed in every tissue of the body, particularly in nervous tissue such as brain. They occur particularly in the outer leaflet of the plasma membrane, where they contribute to cell surface carbohydrates.

The major glycolipids found in animal tissues are glycosphingolipids. They contain ceramide and one or more sugars. Galactosylceramide is a major glycosphingolipid of brain and other nervous tissue, found in relatively low amounts elsewhere. It contains a number of characteristic C24 fatty acids, eg, cerebronic acid.

Galactosylceramide can be converted to sulfogalactosylceramide (sulfatide), present in high amounts in myelin. Glucosylceramide is the predominant simple glycosphingolipid of extraneural tissues, also occurring in the brain in small amounts.

Gangliosides are complex glycosphingolipids derived from glucosylceramide that contain in addition one or more molecules of sialic acid. Neuraminic acid is the principal sialic acid found in human tissues. Gangliosides are present in nervous systems in high concentration.

### **LIPO PROTEINS OF PLASMA**

In plasma, cholesterol and triglycerides form integral component of macromolecule complex called lipoprotein which are conjugated proteins. Lipid part is the prosthetic group and lipid free protein are designated as apolipoproteins or apo proteins. Protein separation including electrophoresis and ultracentrifugation shows progress in lipoprotein chemistry.

Teselius et al in 1941 reported existence of two lipoprotein classes separated by moving boundary electrophoresis.

In 1954 Gofmen et al separated lipoproteins by ultra centrifugation into five major density classes.

## **OVERVIEW OF LIPOPROTEINS AND LIPOPROTEIN METABOLISM**

Lipoproteins are microemulsions composed of lipids (cholesterol, cholesteryl ester, triglyceride and phospholipid) and proteins (apoproteins). Their function is to transport non water soluble cholesterol and triglycerides in plasma. Lipoproteins are spherical particles containing a central core of non-polar lipids (primarily triglycerides and cholesteryl ester) and a surface monolayer of phospholipids and apoproteins

Lipoproteins have been classified on the basis of their densities during ultracentrifugation into<sup>46</sup>:

- 1) Chylomicrons
- 2) Very low density lipoprotein (VLDL)
- 3) Low density Lipoprotein (LDL)
- 4) High density lipoprotein
- 5) Lipoprotein a

## CLASSIFICATION OF LIPOPROTEINS <sup>46</sup>

| Lipoproteins | Source                                       | Diameter<br>(nm) | Density<br>(g/ml) | Composition    |              | Main lipid<br>components     |
|--------------|--|------------------|-------------------|----------------|--------------|------------------------------|
|              |  |                  |                   | Protein<br>(%) | Lipid<br>(%) |                              |
| Chylomicrons | Intestine                                    | 90-1000          | <0.95             | 1-2            | 98-99        | Triacylglycerol              |
| VLDL         | Liver  | 30-90            | 0.95-1.006        | 6-8            | 92-94        | Triacylglycerol              |
| LDL          | VLDL   | 20-25            | 1.019-1.063       | 21             | 79           | Cholesterol                  |
| Lp(a)        |  | 25-35            | 1.040-1.063       | 36             | 55           | Cholesterol                  |
| HDL          | Liver,<br>intestine,<br>VLDL<br>chylomicrons |                  |                   |                |              | Phospholipids<br>Cholesterol |
| HDL1         |  | 20-25            | 1.019-1.063       | 32             | 68           |                              |
| HDL2         |  | 10-20            | 1.063-1.125       | 33             | 67           |                              |
| HDL3         |  | 5-10             | 1.125-1.210       | 57             | 43           |                              |

## **CHYLOMICRONS**

Chylomicrons are the largest of lipoproteins. They measure about 90-1000nm in diameter and they have the least density as compared to other classes of lipoproteins.

These are particles that are primarily triglyceride bearing and are produced by the intestine after exogenous fat undergoes digestion. These are responsible for the transport of dietary triglycerides and cholesterol. Dietary triglycerides are hydrolyzed in the gut, releasing monoglycerides and fatty acids that are then reesterified to form triglycerides in the intestinal mucosal cell.

These triglycerides are assembled with newly absorbed cholesterol, apoprotein B48 and the A apoproteins. Upon secretion from the enterocyte, these assembled particles enter the lymphatic circulation and then the bloodstream, where they acquire C apoproteins and apo E by transfer from HDL.

As chylomicrons enter the plasma, the triglycerides are rapidly hydrolyzed by the enzyme lipoprotein lipase (LPL), which resides on the surface of capillary endothelial cells. LPL is synthesized primarily in adipose tissue and striated muscle. It is secreted and transported to the endothelial surface, where it acts on triglyceride rich particles. Its action requires the presence of apo CII on the surface of the lipoprotein, whereas apo CIII inhibits LPL.

As triglyceride is depleted from the chylomicrons, phospholipids and A and C apoproteins are transferred to HDL. The residual chylomicron particle, which has lost 80 to 90 % of its triglyceride and is now relatively cholesterol enriched is called chylomicron remnant.

## **VERY LOW DENSITY LIPOPROTEINS**

VLDLs are synthesized in the endoplasmic reticulum of hepatocytes and are composed of endogenous triglyceride derived from plasma free fatty acids, chylomicron remnants and from de novo lipogenesis.

Nascent VLDLs are secreted into the circulation; contain apoB100 and small amounts of apo C and apo E. After VLDLs enter the circulation, they are metabolized in the same manner as chylomicrons by the enzyme LPL, with the fatty acids that are liberated following the same fate as those liberated from chylomicrons. After secretion, VLDLs acquire more C and E apoproteins by transfer from HDL. In addition, free cholesterol is progressively exchanged to HDL, where it is esterified and the cholesteryl ester is returned to VLDL.

As VLDLs become progressively depleted of triglyceride, a portion of the surface, including cholesterol, apolipoproteins C and E and phospholipids is removed and contributes to nascent HDL particles.

## **LOW DENSITY LIPOPROTEINS**

LDLs are products of the metabolism of VLDL. They have got a diameter of 18-25nm. The only apoprotein in LDL is apo B and only one molecule of apo B is present per particle of LDL. Clearance of LDL is mediated by a specific receptor present on the surface of both liver and peripheral cells. Once it is bound to the receptor, the lipoprotein is internalized by an endocytic process. The vesicle then fuses with a lysosome, where enzymes degrade the apoB and hydrolyze the cholesteryl ester to free cholesterol.

The smaller remnants of VLDL are triglyceride depleted, cholesterol rich particles, some of which are isolated in the IDL compartment, although some remain in the VLDL compartment.

VLDL particles are believed to be secreted in a spectrum of sizes with various degrees of triglyceride enrichment. The larger VLDL particles appear to be more rapidly cleared and less likely to be converted to LDL. On the other hand, smaller VLDL particles that are richer in cholesterol may be preferentially converted to LDL.

All peripheral cells express the LDL-receptor (LDL-R), and recycle it to the cell surface upon need for cholesterol. Cholesterol is delivered to these cells through binding of LDL to LDL-R, which triggers endocytosis (internalization) of both species. When the need for cholesterol is satisfied, the recycling of LDL-R is discontinued. Normally, an LDL particle stays in circulation for no more than a few days before being consumed by a cholesterol needing cell.

However, under conditions of sustained cholesterol excess, the particle stays in circulation for longer periods of time, and becomes more vulnerable to undesired modifications (e.g. oxidation). As high levels of oxidized LDL are commonly found in atherosclerotic plaques, they are thought to be the major inducer of atherosclerotic lesions. Hence, LDL became known as bad cholesterol

## **HIGH DENSITY LIPOPROTEINS**

HDLs also are represented by a spectrum of particles of various sizes and densities. HDLs are 18-25nm in diameter.

HDL is synthesized in the liver and intestine as a nascent, discoid-shaped particle that contains predominantly apoA-I, and some phospholipids.<sup>46</sup> Upon maturation, HDL assumes a spherical shape, and the composition of its core lipids becomes very similar to that of LDL. However, the relative higher protein content in HDL renders the particle denser and more resistant to undesired modifications.

Unlike LDL, HDL is not recognized by LDL-R, and cannot deliver cholesterol to tissue cells. Instead, it has the ability to remove excess peripheral cholesterol and return it to the liver for recycling and excretion. This process, called reverse cholesterol transport, is thought to protect against atherosclerosis. Observational studies over the last 2 decades have consistently shown strong correlation between elevated HDL levels and low incidents of coronary artery disease (CAD). Hence HDL has been dubbed “good” cholesterol.

### **LIPOPROTEIN (a)**

Lipoprotein (a) or Lp (a) has been established as an independent CAD risk factor. The structure is similar to that of an LDL molecule linked by a disulphide bridge to apoprotein A.

Lp (a) levels range from 1-100mg /dl with the largest number of values below 20 mg/dl.

Although Lp (a) is structurally similar to LDL, the former appears to be regulated independently and carries an independent relation to overall coronary risk. If serum levels of both LDL and Lp (a) are elevated the risk of CAD is markedly increased.

The mechanism by which high levels of Lp (a) are related to coronary atherosclerosis is unclear. It has been suggested that because of the structural similarities of Lp (a) to plasminogen, high levels of Lp (a) may inhibit the thrombolytic activity of naturally occurring tissue plasminogen activity.

An alternative explanation for the association between elevated Lp (a) levels and atherosclerosis is that Lp (a) may somehow alter the LDL mediated delivery of cholesterol to the atherosclerotic plaque.

## **APO PROTEINS**

Apo proteins are key lipoprotein components that serve both as enzymatic cofactors and as recognition elements that bind to specific receptors on peripheral tissues, including the vascular endothelial cells.

It is the Apo E component of the chylomicron remnant that is recognized by receptors on the hepatocyte. These apoproteins are distinguished alphabetically and numerically as Apo A I through Apo E.

A great deal of research has been conducted in the use of apoproteins as CAD markers. Some investigations have found that the concentration of Apo AI and Apo B 100 are better predictors of CAD than are measurements of total plasma lipids or lipoprotein.

### **Apoprotein – A**

Apo AI, the prototype of Apo A, is a major protein in HDL and also is seen in chylomicrons.

Apo AII is a minor constituent of HDL and does not appear to be present in all species. Human Apo AII is of hepatic origin. Apo AII may be an activator of hepatic triglyceride lipase which hydrolyses triglyceride and phospholipid while utilizing HDL2 as its preferred substrate. Apo A IV is synthesized in the gut and is present in HDL, chylomicrons and as a free protein. It may also be an activator of LCAT.

The genetic codes for Apos AI, A IV and CIII are close together on the long arm of chromosome II. Combined AI CIII deficiency is associated with severe premature atherosclerosis.

### **Apoprotein - B**

Apo B occurs in two forms namely Apo B 100 and B28. Apo B 100 is found in VLDL, IDL, and LDL. Apo B 100 is the primary apoprotein of LDL and accounts for 25% of its weight. It is also the recognition site for the LDL, or Apo B/E receptor on cell surfaces. The gene for Apo B 100 has been localized to chromosome 2.

The structure in the amino acid sequence of human Apo B 100 and the corresponding CDNA messenger CID has recently been determined. A unique editing mechanism introduces a stop codon into the mRNA for Apo B by means of single base change. This allows the biosynthesis of two proteins from a single gene and mRNA with either Apo B -100 or Apo B 48 being synthesized. Apo B 48 is synthesized by small intestine and Apo B 100 is secreted by the liver.

### **Apoprotein - E**

Apo E accounts for about 15 percent of the protein content of VLDL, 7 percent of chylomicron remnants and 2 percent of protein content of HDL.

It can be recognized by the LDL or Apo B/E receptor and by specific Apo E receptors in the liver whose function appears to be the removal of chylomicron remnants Apo E is polymorphic and contains three major alleles, Apo E2, E3 and E4. The various combinations results in homozygote are Apo E2/2 EG/4. Also Apos E2/3, E2/4 and E3/4 exist in the heterozygous state. The polymorphism of Apo E has been determined on a molecular basis and results from the substitution of an amino acid at residues 112 and 158 in the protein. Apo E isoform may account for as much as 15 percent of variability of cholesterol and LDL levels in the population.

Recent Finnish studies suggest that Apo E4 may be associated with increased cholesterol absorption in the GI tract.

**ATP 3 guidelines of Dyslipidemia:**

- Total Cholesterol >200 mg/dl
- Triglycerides >200 mg/dl
- HDL <40 mg/dl
- LDL >190 mg/dl

## **MATERIALS AND METHODS**

### **SOURCE OF DATA**

The material for the present study will be collected from patients who attend the Outpatient department and Inpatient department in BLDEU'S Shri. B. M. Patil Medical College Hospital and Research Centre, Vijayapur from January 2015 to June 2016. The sample size is 60.

### **METHOD OF COLLECTION OF DATA**

- a. By detail history
- b. By detail examination
- c. By relevant investigations like CBC, RBS, Blood urea, Serum creatinine, Lipid profile, Urine routine, Chest X-ray, ECG, CT/ MRI brain as appropriate, Serum uric acid.

### **INCLUSION CRITERIA**

All patients with Ischaemic cerebrovascular accident identified based on clinical as well as laboratory and radiological evaluation (including CT/MRI) admitted in our hospital.

### **EXCLUSION CRITERIA**

- a. Age < 18 years
- b. Patients with chronic intake of hyperuricemic drugs
- c. Patients with conditions which alter serum uric acid levels (lymphoproliferative diseases, polycythemia, myeloproliferative disorders, diabetic ketoacidosis, lactic acidosis)

### **SAMPLE SIZE:**

- With prevalence rate of stroke in India is 1.54/1000 population.<sup>54</sup>
- The sample size is 60, Using formula :

$$n = \frac{Z^2 \cdot P \cdot (100-P)}{d^2}$$

### **Blood sampling and Laboratory procedure for Uric Acid and Lipids**

About 3 ml of blood of venous blood was collected in a clean dry test tube and transported to the laboratory. Serum will be separated from blood within 30 minutes. 0.025 ml of serum is mixed with 1 ml of reagent (peroxidase, uricase, ascorbate oxidase, chromogen). The assay mixture was incubated for 5 minutes at 37°C. After completion of the incubation assay mixture will be measured against reagent blank at 510nm. The blood samples were drawn in the fasting state. The venepuncture was done in the cubital fossa. Tourniquet was used but was released just before sampling to avoid artificial increase in the concentration of serum lipids. About 10 ml of blood was drawn using perfectly dry and sterile syringes and the blood was transferred to dried glass vials. Serum was separated within 2 hrs of collection to prevent artificial changes in concentration of HDL. The blood was centrifuged at 5000 rpm for 10 minutes. The supernatant clear serum was then pipetted out using dry piston pipettes with disposable tips and stored in dry thin walled vials at 4cc. The samples were analyzed the same day. Care was taken to exclude the haemolysed serum.

### **STATISTICAL ANALYSIS**

All characteristics were summarized descriptively. Chi-square ( $\chi^2$ )/Fisher exact test was employed to determine the significance of differences between groups for categorical data. The difference of the means of analysis variables was tested with the unpaired t-test. If the p-value was < 0.05, then the results will be considered to be significant. Data were analyzed using SPSS software v.23.0.


## RESULTS

In present study, 60 patients of ischaemic cerebrovascular accident are considered.

**Table 1 : Distribution of cases according to Sex**

| Sex    | N  | %    |
|--------|----|------|
| Male   | 43 | 71.7 |
| Female | 17 | 28.3 |
| Total  | 60 | 100  |

**Figure 1 : Distribution of cases according to Sex**




**Table 2 : Distribution of cases according to Age (Yrs)**

| Age(Yrs) | N  | %    |
|----------|----|------|
| 25-40    | 6  | 10   |
| 41-55    | 10 | 16.7 |
| 56-70    | 32 | 53.3 |
| 71-85    | 10 | 16.7 |
| >85      | 2  | 3.3  |
| Total    | 60 | 100  |


**Table 3 : Mean Age by Sex**

| Sex    | Age (Yrs) |      |      | p value        |
|--------|-----------|------|------|----------------|
|        | Range     | Mean | SD   |                |
| Male   | 25-85     | 60.1 | 14.2 | 0.009<br>(Sig) |
| Female | 40-93     | 71.0 | 13.8 |                |
| Total  | 25-93     | 63.2 | 14.8 |                |

**Figure 2 : Distribution of cases according to Age (Yrs)**




**Figure 3 : Mean Age by Sex**



**Table 4 : Distribution of cases according to Deficit**

| Deficit         | N  | %    |
|-----------------|----|------|
| Lt. Hemiparesis | 29 | 48.3 |
| Rt. Hemiparesis | 31 | 51.7 |
| Total           | 60 | 100  |


**Figure 4 : Distribution of cases according to Deficit**



**Table 5 : Distribution of cases according to Hyperuricemia**

| <b>Hyperuricemia</b> | <b>N</b> | <b>%</b> |
|----------------------|----------|----------|
| Present              | 18       | 30       |
| Absent               | 42       | 70       |
| Total                | 60       | 100      |


**Figure 5 : Distribution of cases according to Hyperuricemia**



**Table 6 : Mean Distribution of Serum Uric Acid according to Sex**

| Sex    | Serum Uric Acid |      |     | p value    |
|--------|-----------------|------|-----|------------|
|        | Range           | Mean | SD  |            |
| Male   | 3-10            | 5.2  | 1.6 | 0.047(Sig) |
| Female | 2.3-9.9         | 6.2  | 1.9 |            |
| Total  | 2.3-10          | 5.5  | 1.7 |            |


**Figure 6 : Mean Distribution of Serum Uric Acid according to Sex**



**Table 7 : Hyperuricemia and Sex**

| Sex    | Hyperuricemia Present |        | Hyperuricemia Absent |        | Total | p value      |
|--------|-----------------------|--------|----------------------|--------|-------|--------------|
|        | N                     | %      | N                    | %      | N     |              |
| Male   | 7                     | 38.9%  | 36                   | 85.7%  | 43    | <0.001 (Sig) |
| Female | 11                    | 61.1%  | 6                    | 14.3%  | 17    |              |
| Total  | 18                    | 100.0% | 42                   | 100.0% | 60    |              |


**Figure 7 : Hyperuricemia and Sex**



**Table 8 : Mean Distribution of Serum Uric Acid according to Age (Yrs)**

| Age(Yrs) | Serum Uric Acid |      |     | p value |
|----------|-----------------|------|-----|---------|
|          | Range           | Mean | SD  |         |
| 25-40    | 3.8-6.7         | 5.1  | 1.2 | 0.077   |
| 41-55    | 3.2-6.7         | 4.8  | 1.3 |         |
| 56-70    | 2.3-8.5         | 5.4  | 1.6 |         |
| 71-85    | 3.1-10          | 6.2  | 2.2 |         |
| >85      | 6.5-9.9         | 8.2  | 2.4 |         |
| Total    | 2.3-10          | 5.5  | 1.7 |         |


**Figure 8 : Mean Distribution of Serum Uric Acid according to Age (Yrs)**



**Table 9 : Hyperuricemia and Age (Yrs)**

| Age(Yrs) | Hyperuricemia Present |        | Hyperuricemia Absent |        | p value     |
|----------|-----------------------|--------|----------------------|--------|-------------|
|          | N                     | %      | N                    | %      |             |
| 25-40    | 1                     | 5.6%   | 5                    | 11.9%  | 0.023 (Sig) |
| 41-55    | 0                     | 0.0%   | 10                   | 23.8%  |             |
| 56-70    | 10                    | 55.6%  | 22                   | 52.4%  |             |
| 71-85    | 5                     | 27.8%  | 5                    | 11.9%  |             |
| >85      | 2                     | 11.1%  | 0                    | 0.0%   |             |
| Total    | 18                    | 100.0% | 42                   | 100.0% |             |

**Figure 9 : Hyperuricemia and Age (Yrs)**




**Table 10 : Correlation of Serum Uric Acid with Selected parameters**

| <b>Correlation of Serum Uric Acid with following parameters</b> | <b>correlation coeff (r)</b> | <b>p value</b> |
|---|------------------------------|----------------|
| Age   | 0.242                        | 0.062          |
| TLC   | 0.155                        | 0.236          |
| RBS   | -0.016                       | 0.9            |
| BU  | 0.557                        | 0.001 (sig)    |
| SC  | 0.336                        | 0.009 (sig)    |
| TC  | -0.159                       | 0.225          |
| TG  | 0.14                         | 0.287          |
| HDL   | -0.143                       | 0.276          |
| LDL   | -0.114                       | 0.384          |
| VLDL  | -0.096                       | 0.465          |

**Table 11 : Distribution of cases according to Lipid profile**

| <b>Lipid profile</b> | <b>N</b> | <b>%</b> |
|----------------------|----------|----------|
| Dyslipidemia         | 49       | 81.7     |
| Normal               | 11       | 18.3     |
| Total                | 60       | 100.0    |


**Figure 10 : Distribution of cases according to Lipid profile**



**Table 12 : Dyslipidemia and Age**

| Age(Yrs) | Dyslipidemia |        | Normal |        | p value |
|----------|--------------|--------|--------|--------|---------|
|          | N            | %      | N      | %      |         |
| 25-40    | 4            | 8.2%   | 2      | 18.2%  | 0.513   |
| 41-55    | 8            | 16.3%  | 2      | 18.2%  |         |
| 56-70    | 29           | 59.2%  | 3      | 27.3%  |         |
| 71-85    | 7            | 14.3%  | 3      | 27.3%  |         |
| >85      | 1            | 2.0%   | 1      | 9.1%   |         |
| Total    | 49           | 100.0% | 11     | 100.0% |         |


**Figure 11 : Dyslipidemia and Age**



**Table 13 : Dyslipidemia and Sex**

| Sex    | Dyslipidemia |        | Normal |        | p value |
|--------|--------------|--------|--------|--------|---------|
|        | N            | %      | N      | %      |         |
| Male   | 36           | 73.5%  | 7      | 63.6%  | 0.299   |
| Female | 13           | 26.5%  | 4      | 36.4%  |         |
| Total  | 49           | 100.0% | 11     | 100.0% |         |


**Figure 12 : Dyslipidemia and Sex**



**Table 14 : Patient having both Hyperuricemia and Dyslipidemia**

| <b>Hyperuricemia &amp; Dyslipidemia</b> | <b>N</b> | <b>% (out of 60)</b> |
|---|----------|----------------------|
| Male                                    | 6        | 10.0%                |
| Female                                  | 9        | 15.0%                |
| Total                                   | 15       | 25.0%                |


**Figure 13 : Patient having both Hyperuricemia and Dyslipidemia**



**Table 15 : Abnormal Lipid Profile and Hyperuricemia**

| Lipid Profile  | Hyperuricemia |       |        |       |       | p value     |
|----------------|---------------|-------|--------|-------|-------|-------------|
|                | Present       |       | Absent |       | Total |             |
|                | N             | %     | N      | %     | N     |             |
| Increased TC   | 3             | 16.7% | 15     | 83.3% | 18    | 0.140       |
| Increased TG   | 2             | 40.0% | 3      | 60.0% | 5     | 0.610       |
| Decreased HDL  | 16            | 37.2% | 27     | 62.8% | 43    | 0.049 (Sig) |
| Increased LDL  | 2             | 50.0% | 2      | 50.0% | 4     | 0.366       |
| Increased VLDL | 2             | 33.3% | 4      | 66.7% | 6     | 0.851       |


**Figure 14 : Abnormal Lipid Profile and Hyperuricemia**



**Table 16 : Abnormal Lipid Profile and Sex**

| Lipid Profile  | Male |       | Female |       | Total | p value |
|----------------|------|-------|--------|-------|-------|---------|
|                | N    | %     | N      | %     | N     |         |
| Increased TC   | 13   | 72.2% | 5      | 27.8% | 18    | 0.95    |
| Increased TG   | 3    | 60.0% | 2      | 40.0% | 5     | 0.545   |
| Decreased HDL  | 31   | 72.1% | 12     | 27.9% | 43    | 0.907   |
| Increased LDL  | 2    | 50.0% | 2      | 50.0% | 4     | 0.320   |
| Increased VLDL | 4    | 66.7% | 2      | 33.3% | 6     | 0.774   |


**Figure 15 : Abnormal Lipid Profile and Sex**



**Table 17 : Abnormal Lipid Profile and Age(Yrs)**

| Age(Yrs)       | 25-40 |      | 41-55 |       | 56-70 |       | 71-85 |       | >85 |      | Total<br>N | P<br>value |
|----------------|-------|------|-------|-------|-------|-------|-------|-------|-----|------|------------|------------|
|                | N     | %    | N     | %     | N     | %     | N     | %     | N   | %    |            |            |
| Increased TC   | 0     | 0.0% | 3     | 16.7% | 4     | 77.8% | 1     | 5.6%  | 0   | 0.0% | 18         | 0.084      |
| Increased TG   | 0     | 0.0% | 1     | 20.0% | 2     | 40.0% | 2     | 40.0% | 0   | 0.0% | 5          | 0.604      |
| Decreased HDL  | 3     | 7.0% | 8     | 18.6% | 2     | 53.5% | 7     | 16.3% | 2   | 4.7% | 43         | 0.639      |
| Increased LDL  | 0     | 0.0% | 1     | 25.0% | 3     | 75.0% | 0     | 0.0%  | 0   | 0.0% | 4          | 0.765      |
| Increased VLDL | 0     | 0.0% | 1     | 16.7% | 2     | 33.3% | 3     | 50.0% | 0   | 0.0% | 6          | 0.212      |

**Figure 16 : Abnormal Lipid Profile and Age(Yrs)**



## DISCUSSION

Stroke continues to have a great impact on public health. Stroke is frequent, recurring, and is more often disabling than fatal. Although some determinants of stroke, such as age, gender, race, ethnicity and heredity cannot be modified, they are risk markers. However controlling the more important modifiable factors like serum uric acid and lipid levels may reduce the incidence of the disease.<sup>47</sup>

In our study a total of 60 patients with ischemic stroke were included. Out of which 43 (71.7%) were males and 17 (28.3%) were females. The mean age of the patients was  $63.2 \pm 14.8$ . Mean serum uric acid levels in the patients studied was  $5.5 \pm 1.7$ , and 18 patients (30%) were hyperuricemic. Serum uric acid levels were significantly higher in females ( $6.2 \pm 1.7$ ) compared to the males ( $5.2 \pm 1.6$ ).

A study by Mehrpour et al.<sup>48</sup> was done in 55 patients with acute ischaemic stroke. Of which, 25 of were females and 30 were males. The mean age of the patients was  $67 \pm 14$  years. Mean serum uric acid level was  $5.94 \pm 1.70$  mg/dl. 47.3% of patients were hyperuricemic. Uric acid levels were significantly higher in males than females. Hyperuricemia was associated with increase in levels of triglycerides and LDL-C.

Bhadra J et al.<sup>49</sup> evaluated 38 patients with mean age  $59.28 \pm 12.31$ . Serum uric acid levels were significantly higher in study subjects and statistically significant correlation was seen with TG and VLDL and inverse association with HDL in the cases.

Study by Biyani VV et al.<sup>50</sup> studied 100 patients among which 68 were males and 32 females. The patients with hyperuricemia were mostly in the age group of 60-69 years. 49% of the patients had hyperuricemia. Similar results were concluded in studies by Millinois et al and Patil T et al.

In the present study dyslipidemia was seen in 49 (81.7%) patients (Males were 36 and females were 13). It was predominant in the age group between 56-70 years. More than one lipid parameter was seen to be deranged in 10 male patients and 4 female patients. Of them 87.75% of patients had low HDL levels, 36.73%, 10.20% and 8.16% patient had high cholesterol, triglycerides and LDL levels respectively. The study also showed that 15 patients had both hyperuricemia and dyslipidemia.

Baluch U et al.<sup>51</sup> studied 53 patients of which 32 were males and 21 were females. 28% of patients were in age group of 61-70 years. 19% patient had dyslipidemia of them, 18% had low HDL, while high LDL, cholesterol and triglycerides were observed in 26%, 24% and 32% respectively.

Albucher J.F. et al.<sup>52</sup> concluded that low HDL cholesterol was the only serum lipid index to be associated to an increased risk of stroke in the population as seen in our study.

Sarmah D et al.<sup>53</sup> proved a significant correlation between uric acid and TC, TG and LDL, and a significant negative correlation between uric acid and HDL. The similar findings were seen in our study.

## CONCLUSION

Developing countries like India are facing a double burden of communicable and non-communicable diseases. Stroke is one of the leading causes of death and disability in India. Stroke is becoming an important cause of premature death and disability in low-income and middle-income countries like India, largely driven by demographic changes and enhanced by the increasing prevalence of the key modifiable risk factors like hyperuricemia and dyslipidemia. Both of these factors are involved in atherosclerosis and endothelial dysfunction which leads to narrowing of vessels and leads to ischaemia.

Following are the important findings observed in our study:

- 30% of the patients were hyperuricemic with the mean age of  $63.2 \pm 14.8$ . Mean serum uric acid levels was  $5.5 \pm 1.7$ , these findings were similar to the earlier studies. However, it was noted that hyperuricemia was higher in females compared to the males in comparison to available data.
- Statistically significant number of patients showed dyslipidemia, which was common in males in age group of 56-70 years. Of them 87.75% of patients had low HDL levels, 36.73%, 10.20% and 8.16% patient had high cholesterol, triglycerides and LDL levels respectively. More than one lipid parameter was deranged in 28.5% of patients. The study also showed that 25% patients had both hyperuricemia and dyslipidemia.

- HDL-C levels were significantly low and total cholesterol, triglyceride and LDL-C levels was high among the study patients.

In conclusion hyperuricemia and dyslipidemia can be taken as risk factors for ischaemic cerebrovascular accidents.

## SUMMARY

A hospital based study to determine “Serum uric acid levels and serum lipid levels in patients with ischaemic cerebrovascular accident” was conducted during the period of January 2015 to June 2016. All patients with Ischaemic cerebrovascular accident identified based on clinical, laboratory and radiological evaluation (including CT/MRI) admitted in our in-patient department were included in the study. All patients were subjected to detail history, examination and relevant investigations like CBC, RBS, Serum uric acid, Blood urea, Serum creatinine, Lipid profile, Urine routine, Chest X-ray, ECG, CT/ MRI brain as appropriate,.

Following are the salient findings of the study:

Significant increase in serum uric acid, total cholesterol, triglyceride, LDL-C and decreased levels of HDL-C were observed in our study patients.

This study signifies that hyperuricemia and dyslipidemia can lead to ischaemic stroke and they both are modifiable risk factors.

## BIBLIOGRAPHY

1. Smith W.S, English J.D, Johnston S.C. Cerebrovascular diseases. In: Longo D.N, Kasper D.L, Jameson J.L, Fauci A.S, Hauser S.L, Loscalzo J, editors. *Harrisons principles of internal medicine*, 18th edn. USA: Mc graw hill medical; 2012.p.-3270.
2. Goldstein L.B. Ischemic cerebrovascular disease. *Goldman-Cecil Medicine*, 25th edn. USA: Elsevier Saunders; 2016.p.-2434.
3. Weir J.C, Muir S.C, Walters R. M, Lees K,R. Serum urate as an independent predictor of poor outcome and future vascular events after acute stroke.*Stroke*.2003;34:1951-6.
4. Bansal B.C, Gupta R.R, Bansal M.R, Prakash C. Serum lipids and uric acid relationship in ischemic thrombotic cerebrovascular disease. *Stroke* 1975;6:304-7.
5. Hariklia VD, Apostolos H, Haralambosk. The Role of Uric Acid in Stroke. The Issue Remains Unresolved. *The Neurologist*. 2008; 14: 238-242.
6. Feigin VL, Lawes CM, Bennett DA, et al. Worldwide stroke incidence and early case fatality reported in 56 population- based studies: a systematic review. *Lancet Neurol*. Llibre J, valhuerdi A, Fernandez O. prevalence of stroke and Associated Risk factors in Older Adults in Havana city and Matanzas Provinus, Cuba (10/66 Population- Based study). *MEDICC Review*. 2010; 12: 20-24.
7. Llibre J, valhuerdi A, Fernandez O. prevalence of stroke and Associated Risk factors in Older Adults in Havana city and Matanzas Provinus, Cuba (10/66 Population- Based study). *MEDICC Review*. 2010; 12: 20-24.

8. Barr WG. Uric acid. In: Walker HK, Hall WD, Hurst JW, editors. *Clinical Methods. The History, Physical, and Laboratory Examination*, 3rd ed. Boston: Butterworth; 1990. P: 760-763.
9. Heo SH, Lee SH. High levels of serum Uric acid are associated with silent brain infarction. *Journal of the Neurological Sciences*. 2010; 297:6-15.
10. Berry C, Hamilton CA, Brosnan MJ, Magill FG, Berg GA, McMurray JJ, Dominiczak AF. Investigation into the sources of superoxide in human blood vessels: Angiotensin II increases superoxide production in human internal mammary arteries. *Circulation*. 2000;13:2206–2212.
11. DeScheerder IK, van de Kraay AM, Lamers JM, Koster JF, deJong JW, Serruys PW. Myocardial malondialdehyde and uric acid release after short-lasting coronary occlusions during angioplasty: potential mechanisms for free radical generation. *Am J Cardiol*. 1991; 68: 392–395.
12. Kang DH, Park SK, Lee IK, Johnson RJ. Uric acid-induced C-reactive protein expression: implication on cell proliferation and nitric oxide production of human vascular cells. *J Am Soc Nephrol*. 2005;13:3553–3562.
13. Ginsberg MH, Kozin F, O'Malley M, McCarty DJ. Release of platelet constituents by monosodium urate crystals. *J Clin Invest*. 1997; 60: 999–1007.
14. Netea MG, Kullberg BJ, Block WL, Netea RT, van der Meer JW. The role of hyperuricemia in the increased cytokine production after lipopolysaccharide challenge in neutropenic mice. *Blood*. 1997;89:577–582.
15. Mazzali M, Hughes J, Kim YG, Jefferson JA, Kang D-H, Gordon KL, Lan HY, Kivlighn S, Johnson RJ. Elevated uric acid increases blood pressure in

- the rat by a novel crystal-independent mechanism. *Hypertension*. 2001; 38: 1101–1106.
16. Watanabe S, Kang D-H, Feng L, Nakagawa T, Kanellis J, Lan H, Mazzali M, Johnson RJ. Uric acid, hominoid evolution, and the pathogenesis of salt-sensitivity. *Hypertension*. 2002; 40: 356–360.
  17. Sanchez-Lozada LG, Tapia E, Avila-Casado C, Soto V, Franco M, Santamaria J, Nakagawa T, Rodriguez-Iturbe B, Johnson RJ, Herrera-Acosta J. Mild hyperuricemia induces glomerular hypertension in normal rats. *Am J Physiol Renal Physiol*. 2002; 283: F1105–F1110.
  18. Kang DH, Nakagawa T, Feng L, Watanabe S, Han L, Mazzali M, Truong L, Harris R, Johnson RJ. A role for uric acid in the progression of renal disease. *J Am Soc Nephrol*. 2002; 13: 2888–2897.
  19. Mazzali M, Kanellis J, Han L, Feng L, Xia Y-Y, Chen Q, Kang D-H, Gordon KL, Watanabe S, Nakagawa T, et al. Hyperuricemia induces a primary renal arteriopathy in rats by a blood pressure-independent mechanism. *Am J Physiol Renal Physiol*. 2002; 282: F991–F997.
  20. Suarna C, Dean RT, May J, Stocker R. Human atherosclerotic plaque contains both oxidized lipids and relatively large amounts of alpha-tocopherol and ascorbate. *Arterioscler Thromb Vasc Biol*. 1995; 15: 1616–1624.
  21. Vannorsdall TD, Jinnah HA, Gordon B, Kraut M, Schretlen DJ. Cerebral ischemia mediates the effect of serum uric acid on cognitive function. *Stroke*. 2008;13:3418–3420.
  22. Dodson PM and Horton RC. 1987: "The nature of atheroma". *Proc. Nutr. Soc*, 46: 331-336.

23. Steinberg D et al 1989: "Beyond cholesterol: modification of low density lipoprotein that increase its atherogenicity". *N. Engl. J Med.* 320: 915-924.
24. Howard BY et al 1983: "VLDL triglyceride metabolism in NIDDM. Relationship of plasma insulin and free fatty acids". *Diabetes*, 32: 271- 276.
25. Scheele KW. Examen Chemicum Calculi Urinarii, Opuscula 11,p,73,1776.Cited from Levene,P.A and Bass,L.W:Nucleic Acids.New York,Chemical catalogcompany,1931.
26. Wollaston,W.H. On Gouty and urinary concertions, Phil. Trans. London,1797,87:386.
27. Becker, Michael A, M.D., Schumaker, H. Ralph Jr., M.D., Wortmann, Robert L. M.D., and others; Febuxostat Compared with Allopurinol in Patients with Hyperuricemia and Gout. *New England Journal of Medicine* 005; 353: 2450-61.
28. Note on the estimation of uric acid by the Hopkins-folin method. By Robert Campbell Garry. Institute of Physiology, Glasgow University.
29. Sir Archibald Garrod and Alkaptonuria –‘story of metabolic genetics’C Prasad1,\* , PA Galbraith2.
30. Alkylation of oxypurines by  $\alpha,\omega$ -dihaloalkanes 1. Reaction of xanthine sodium salts and some N- methyl derivatives with  $\alpha,\omega$ -dihaloalkanesI. Sh. Salikhov, V. O. Kuznetsov and V. S. Reznik.
31. Baillie, J.K; M.G.Bates, A.A. Thompson, W.S. Waring, R.W.Partridge, M.F.chnopp, A. Simpson, F. Gulliver – Sloan, S.R. Maxwell, D.J. Webb(2007-05). Endogenous uric acid production augments plasma antioxidant capacity in healthy lowland subjects exposed to high altitude. *Chest* 131(5): 1473-1478.

32. S.R.J.Maxwell, Thomason, D.Sandler, C.Leguen, M.A.Baxter, G.H.G. Thorpe, A.F. Jones, A.H. Barnett. ‘ ‘ Antioxidant status in patients with uncomplicated insulin dependent and non insulin dependent diabetes mellitus’’. European Journal of clinical investigation(1997)27,484-490.
33. Doring A, Gieger C, Mehta D, et al.(April 2008). ‘ ‘SLC2A9 influences uric acid concentrations with pronounced sex specific effects’’. Nature Genetics 40 (4);430-6.
34. Best and Taylor, Physiological Basis of Medical Practice ‘ ‘Uric Acid’’, 12th edition, Williams and Wilkins, Section 4 30:479.
35. E medicine Feb. 9 2009 Uric Acid Nephropathy. Author: Mark T Fahlen, MD, Inc Coauthor(s): Mahendra Agraharkar, MD, MBBS, FACP, FASN, Clinical Associate Professor of Medicine, Baylor College of Medicine, feb 9 2009.
36. Identification of the causes of persistent hyperuricaemia. The Lancet, Volume 337, Issue 8755, Pages 1461-1463 B.Emmerson.
37. Hypouricemia in the Syndrome of Inappropriate Secretion of Antidiuretic Hormone Laurence H. Beck, M.D. N Engl J Med 1979; 301:528-530 September 6, 1979 new England journal of medicine.
38. Abnormalities in uric acid metabolism: concept and classification. Nippon Rinsho 1996 Dec; 54(12):3243-7 Akaoka I, Kamatani N 2nd Department of Internal Medicine, Teikyo University School of Medicine.
39. Hypouricaemia and increased renal uric acid clearance associated with hyperparathyroidism. T Gibson, H P Sims, and S A Jimenez Ann Rheum Dis. 1976 August; 35(4): 372–376.

40. Determination of uric acid levels in uraemia by enzymatic and colorimetric techniques P. J. N. Howorth, Joan F. Zilva *J Clin Pathol* 1968;21:192-195 doi:10.1136/jcp.21.2.192.
41. Update on Gout and Hyperuricemia J. F. Baker; H. Ralph Schumacher Posted: 02/22/2010; *Int J Clin Pract.* 2010;64(3):371-377. © 2010.
42. Hypo-uricemia in the syndrome of inappropriate secretion of antidiuretic hormone: A prospective study Passeron A, Blanchard A, Capron L. *Rev Med Interne.* 2010 Oct; 31(10):665-9. Epub 2010 Jul 3.
43. Glantzounis GK, Tsimoyiannis EC, Kappas AM, Galaris DA (2005). ‘‘Uric acid and oxidative stress’’ *Current Pharmaceutical Design* 11(32); 4145 -51.
44. Uricase based methods for determination of uric acid in serum, Y S Zhao, X Y Yang, W Lu, H Liao, F Liao et al.(2009) Volume:164, Issue:1-2, pages:1-6.
45. Botham KM and Mayes PA. *Harper’s Illustrated Biochemistry.* 28th Edition. Page no 121.
46. Howard BV and Howard WJ. *Pathophysiology and Treatment of Lipid Disorders in Diabetes.* Joslin’s diabetes mellitus. Fourteenth edition, Page no: 563-582.
47. Sacco RL, Benjamin EJ, Broderick JP, et al. American Heart Association Prevention Conference IV: prevention and rehabilitation of stroke: risk factors. *Stroke.*
48. Mehrpour M, Khuzan M, Najimi N, Motamed M.R, Fereshtehnejad S.M. Serum uric acid level in acute stroke patients. *Medical Journal of Islamic Republic of Iran,* Vol. 26, No. 2, May. 2012, pp. 66-72


49. Bhadra J, Mahor DS, Seth S, Ghalaut VS, Dhupper V, Kulshrestha MR. Uric acid in men with acute stroke. *IJIMS*, 2015, vol 2, no.5, 146-152.
50. Biyani VV, Mane RR, Khyalappa R. Association of serum uric acid an independent risk factor in acute ischemic stroke. *Sch. J. App. Med. Sci.*, 2016; 4(5B):1539-1543.
51. Baluch UT, Kiani I, Nisa ZU, Badshah M. Association of Dyslipidemia and Ischemic Stroke. *Ann. Pak. Inst. Med. Sci.* 2008;4(3): 165-167.
52. Albucher JF et al. Serum lipids in young patients with ischaemic stroke: a case control study. *J Neurol Neurosurg Psychiatry* 2000 69: 29-33.
53. Sarmah D, Sharma B. A correlative study of uric acid with lipid profile. *Asian Journal of Medical Science*, Volume-4(2013).
54. Park K. Non communicable diseases. *Preventive and Social Medicine*, 21<sup>st</sup> edition. Jabalpur: Banarsidas Bhanot; 2011.p.-349.

## ANNEXURES

### ETHICAL CLEARANCE



B.L.D.E. UNIVERSITY'S  
SHRI.B.M.PATIL MEDICAL COLLEGE, BIJAPUR-586 103  
INSTITUTIONAL ETHICAL COMMITTEE



#### **INSTITUTIONAL ETHICAL CLEARANCE CERTIFICATE**

The Ethical Committee of this college met on 22-11-2014 at 3:30pm to scrutinize the Synopsis of Postgraduate Students of this college from Ethical Clearance point of view. After scrutiny the following original/corrected & revised version synopsis of the Thesis has been accorded Ethical Clearance.

Title: "Serum Uric Acid Levels and Serum Lipid Levels in Patients with Ischaemic cerebrovascular Accident"  
— x — x — x — x —

Name of P.G. student Dr. Tavish Arora.

Dept of General Medicine

Name of Guide/Co-investigator Dr. P. G. Mantur. Asso. Professor.

Dept of General Medicine

for   
DR. TEJASWINI VALLABHA  
CHAIRMAN  
INSTITUTIONAL ETHICAL COMMITTEE  
BLDEU'S, SHRI.B.M.PATIL  
MEDICAL COLLEGE, BIJAPUR.

Following documents were placed before E.C. for Scrutinization

- 1) Copy of Synopsis/Research project.
- 2) Copy of informed consent form
- 3) Any other relevant documents.

## **CONSENT FORM**

I ..... exercising free power of choice hereby give my written consent to be included as a subject in the study **“SERUM URIC ACID LEVELS AND SERUM LIPID LEVELS IN PATIENTS WITH ISCHAEMIC CEREBROVASCULAR ACCIDENT”** conducted by **Dr. TAVISH ARORA** and undergo the necessary investigations required for this study and I fully consent for the same.

I am over 18 yrs of age and I have been explained to my satisfaction by the attending Physician, in the language I understand, about the purpose of the study .I have also understood that the investigator will maintain confidentiality regarding my identity.

**SIGNATURE OF DOCTOR**

**SIGNATURE OF PATIENT/RELATIVE**

**NAME OF THE DOCTOR**

**NAME OF PATIENT/RELATIVE**

**Dr. TAVISH ARORA**

**RELATIONSHIP (IF RELATIVE)**

**DATE:**

**PROFORMA**

**BLDE'S SHRI B.M.PATIL MEDICAL COLLEGE HOSPITAL**

**AND RESEARCH CENTRE, VIJAYAPUR**

**"SERUM URIC ACID LEVELS AND SERUM LIPID LEVELS IN**

**PATIENTS WITH ISCHAEMIC CEREBROVASCULAR**

**ACCIDENT"**

Patient Name:

I.P number:

Age:

Sex:

Address:

Occupation:

Date of admission:

Date of discharge:

***SYMPTOMS :***

- |                           |        |
|---------------------------|--------|
| 1. Fever                  | Yes/No |
| 2. Preceding delivery     | Yes/No |
| 3. Preceding convulsions  | Yes/No |
| 4. Preceding headache     | Yes/No |
| 5. Unconsciousness        | Yes/No |
| 6. Right sided hemiplegia | Yes/No |
| 7. Left sided hemiplegia  | Yes/No |
| 8. Monoplegia             | Yes/No |

***TREATMENT HISTORY:***

Chronic diuretic use Yes/No

Chronic salicylate use Yes/No

Use of any other drug altering uric acid level-

***PERSONAL HISTORY:***

Habits:

H/o Smoking Yes/No

H/o Alcohol consumption Yes/No

***PAST HISTORY:***

Any medical disease affecting uric acid levels-

***PHYSICAL EXAMINATION:***

GENERAL CONDITION:

Pallor: Yes/No

Icterus: Yes/No

Lymphadenopathy: Yes/No

Cyanosis: Yes/No

Clubbing: Yes/No

Edema: Yes/No

VITALS:

Temperature:

Pulse:

Respiratory rate:

Blood pressure:

Pt condition on discharge:

Improved/ Worsened/Same/Expired

***SYSTEMIC EXAMINATION:***

C.N.S.:

C.V.S.:

P.A.:

R. S:

***LABORATORY INVESTIGATIONS:***

· **CBC**

· **RBS**

· **Blood urea**

· **Serum creatinine**

· **Lipid Profile**

· **Urine routine**

- **ECG**

- **CT/ MRI brain**

- **Serum uric acid**

***DIAGNOSIS:***

## KEY TO MASTER CHART

|      |   |                              |
|------|---|------------------------------|
| TLC  | – | Total leukocyte count        |
| RBS  | – | Random blood sugar           |
| BU   | – | Blood urea                   |
| SC   | – | Serum creatinine             |
| TC   | – | Total cholesterol            |
| TG   | – | Triglycerides                |
| HDL  | – | High density lipoprotein     |
| LDL  | – | Low density lipoprotein      |
| VLDL | – | Very low density lipoprotein |
| Hyp  | – | Hyperuricemia                |

## MASTER CHAR

| Sl. No. | Name                 | Age | Sex | TLC      | RBS   | BU    | SC    | TC    | TG    | HDL   | LDL   | VLDL  | Serum Uric Acid | Deficit         | Lipid profile | Hyp     |
|---------|----------------------|-----|-----|----------|-------|-------|-------|-------|-------|-------|-------|-------|-----------------|-----------------|---------------|---------|
|         |                      |     |     | Cells/mm | mg/dl           |                 |               |         |
| 1       | Mallappa Toravi      | 70  | M   | 9180     | 85    | 55    | 2.8   | 191   | 78    | 47    | 128.4 | 15.6  | 7.5             | Rt. Hemiparesis | Normal        | Present |
| 2       | Shivappa Salytagi    | 36  | M   | 1880     | 83    | 15    | 7     | 170   | 112   | 40    | 107.6 | 22.4  | 3.8             | Lt. Hemiparesis | Normal        | Absent  |
| 3       | Tungabai Golpe       | 70  | F   | 8270     | 96    | 29    | 1.4   | 158   | 85    | 34    | 107   | 17    | 6.0             | Rt. Hemiparesis | Dyslipidemia  | Present |
| 4       | Somaning Gonasagi    | 70  | M   | 21040    | 110   | 54    | 0.8   | 239   | 132   | 49    | 163.6 | 26.4  | 4.6             | Rt. Hemiparesis | Dyslipidemia  | Absent  |
| 5       | Malleshappa Nadigeri | 65  | M   | 6900     | 109   | 21    | 0.9   | 235   | 158   | 37    | 166.4 | 31.6  | 5.4             | Lt. Hemiparesis | Dyslipidemia  | Absent  |
| 6       | Dattatreya Jadhav    | 80  | M   | 7120     | 245   | 168   | 2.3   | 147   | 235   | 22    | 78    | 47    | 10              | Rt. Hemiparesis | Dyslipidemia  | Present |
| 7       | Sidramayya Kammamath | 52  | M   | 6180     | 71    | 20    | 0.9   | 156   | 78    | 43    | 97.4  | 15.6  | 6.7             | Rt. Hemiparesis | Normal        | Absent  |
| 8       | Arjun Jarale         | 42  | M   | 11330    | 102   | 18    | 0.7   | 150   | 147   | 28    | 92.6  | 29.4  | 3.6             | Rt. Hemiparesis | Dyslipidemia  | Absent  |
| 9       | Anasubai Biradar     | 70  | F   | 7790     | 118   | 17    | 0.8   | 178   | 71    | 43    | 120.8 | 14.2  | 6.1             | Lt. Hemiparesis | Normal        | Present |
| 10      | Basavaraj Nagumeti   | 25  | M   | 4820     | 117   | 24    | 0.7   | 144   | 117   | 45    | 75.6  | 23.4  | 4.7             | Lt. Hemiparesis | Normal        | Absent  |
| 11      | Bhimashankar Biradar | 70  | M   | 8120     | 266   | 44    | 1.4   | 232   | 197   | 27    | 165.6 | 39.4  | 4.5             | Rt. Hemiparesis | Dyslipidemia  | Absent  |
| 12      | Ramappa Nidoni       | 70  | M   | 12530    | 141   | 62    | 1.1   | 228   | 136   | 49    | 151.8 | 27.2  | 4.3             | Lt. Hemiparesis | Dyslipidemia  | Absent  |
| 13      | Irannagouda Borwat   | 60  | M   | 6800     | 114   | 30    | 0.7   | 139   | 151   | 32    | 76.8  | 30.2  | 4.4             | Lt. Hemiparesis | Dyslipidemia  | Absent  |

|    |                       |    |   |       |     |    |     |     |     |    |       |      |     |                 |              |         |
|----|-----------------------|----|---|-------|-----|----|-----|-----|-----|----|-------|------|-----|-----------------|--------------|---------|
| 14 | Badusab Chapparaband  | 65 | M | 13960 | 84  | 34 | 0.8 | 158 | 131 | 24 | 107.8 | 26.2 | 3.4 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 15 | Mallamma Medagar      | 80 | F | 7450  | 111 | 42 | 1.7 | 207 | 201 | 29 | 137.8 | 40.2 | 8.5 | Lt. Hemiparesis | Dyslipidemia | Present |
| 16 | Chandbi Attar         | 80 | F | 10790 | 176 | 15 | 0.7 | 180 | 193 | 32 | 109.4 | 38.6 | 5.9 | Rt. Hemiparesis | Dyslipidemia | Present |
| 17 | Basanna Ahanur        | 70 | M | 15650 | 125 | 60 | 1.4 | 248 | 218 | 36 | 168.4 | 43.6 | 6.3 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 18 | Lakshmawwa Kannolli   | 85 | F | 12620 | 116 | 43 | 0.9 | 133 | 82  | 34 | 82.6  | 16.4 | 7.0 | Lt. Hemiparesis | Dyslipidemia | Present |
| 19 | Yallowwa Chalawadi    | 40 | F | 7830  | 95  | 50 | 1   | 129 | 147 | 20 | 90    | 29.4 | 6.7 | Rt. Hemiparesis | Dyslipidemia | Present |
| 20 | Jilani Karjagi        | 65 | M | 8820  | 75  | 17 | 0.7 | 131 | 110 | 30 | 79    | 22   | 4.9 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 21 | Shanabasappa Nivaragi | 61 | M | 9600  | 123 | 12 | 0.6 | 186 | 113 | 36 | 127.4 | 22.6 | 5.0 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 22 | Tarabai Badiger       | 65 | F | 17420 | 150 | 60 | 0.8 | 183 | 160 | 30 | 121   | 32   | 8.0 | Rt. Hemiparesis | Dyslipidemia | Present |
| 23 | Hashmat Khan          | 60 | M | 8200  | 149 | 30 | 1.5 | 208 | 101 | 42 | 145.8 | 20.2 | 6.2 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 24 | Shivalingappa Biradar | 50 | M | 7420  | 92  | 31 | 0.7 | 170 | 177 | 23 | 111.6 | 35.4 | 3.2 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 25 | Parvati Mirji         | 70 | F | 17250 | 148 | 14 | 0.8 | 285 | 117 | 50 | 206.6 | 23.4 | 2.3 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 26 | Gudulal Maniyar       | 60 | M | 9320  | 206 | 16 | 0.8 | 225 | 184 | 35 | 153.2 | 36.8 | 4.0 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 27 | Mayawwa Arjunagi      | 70 | F | 12400 | 110 | 26 | 0.6 | 227 | 209 | 32 | 153.2 | 41.8 | 3.5 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 28 | Basavaraj Naganur     | 45 | M | 7110  | 92  | 12 | 0.8 | 311 | 152 | 24 | 256.6 | 30.4 | 3.9 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 29 | Madiowalappa Jambenal | 60 | M | 10000 | 96  | 36 | 1.2 | 180 | 84  | 24 | 139.2 | 16.8 | 5.1 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 30 | Sidaraya Honakande    | 58 | M | 12220 | 200 | 47 | 2.8 | 174 | 188 | 27 | 109.4 | 37.6 | 5.1 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 31 | Irasangappagouda      | 64 | M | 8080  | 278 | 18 | 0.8 | 130 | 141 | 32 | 69.8  | 28.2 | 3.5 | Rt. Hemiparesis | Dyslipidemia | Absent  |

|    | Biradar             |    |   |       |     |     |     |     |     |    |       |       |     |                 |              |         |
|----|---------------------|----|---|-------|-----|-----|-----|-----|-----|----|-------|-------|-----|-----------------|--------------|---------|
| 32 | Bhimappa Gali       | 40 | M | 6290  | 207 | 20  | 0.8 | 154 | 116 | 22 | 108.8 | 23.2  | 4.0 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 33 | Moula Malur         | 40 | M | 8890  | 113 | 19  | 0.7 | 200 | 173 | 29 | 136.4 | 34.6  | 4.9 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 34 | Malleshi Mantur     | 55 | M | 9280  | 83  | 19  | 0.7 | 120 | 89  | 24 | 78.2  | 17.8  | 3.7 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 35 | Prakash Rathod      | 27 | M | 10800 | 114 | 24  | 1.1 | 144 | 79  | 60 | 68.2  | 15.8  | 6.2 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 36 | Mahadevi Patil      | 93 | F | 22290 | 126 | 147 | 2.1 | 121 | 187 | 22 | 61.6  | 37.4  | 6.5 | Lt. Hemiparesis | Normal       | Present |
| 37 | Shakuntala Channi   | 70 | F | 5450  | 122 | 25  | 1.0 | 172 | 129 | 41 | 105.2 | 25.8  | 4.4 | Lt. Hemiparesis | Normal       | Absent  |
| 38 | Nanasaheb Patil     | 55 | M | 7040  | 110 | 11  | 0.7 | 153 | 144 | 25 | 99.2  | 28.8  | 5.0 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 39 | Vital Chavan        | 57 | M | 6520  | 114 | 17  | 0.7 | 100 | 86  | 23 | 59.8  | 17.2  | 8.5 | Rt. Hemiparesis | Dyslipidemia | Present |
| 40 | Shivkumar Puranik   | 65 | M | 12030 | 94  | 31  | 1.1 | 95  | 106 | 21 | 52.8  | 21.2  | 7.2 | Lt. Hemiparesis | Dyslipidemia | Present |
| 41 | Shakinabee Hundekar | 90 | F | 12640 | 108 | 205 | 5.9 | 88  | 173 | 30 | 90    | 15    | 9.9 | Rt. Hemiparesis | Dyslipidemia | Present |
| 42 | Laxman Sanjeevagal  | 85 | M | 7310  | 110 | 58  | 1.2 | 150 | 114 | 38 | 89.2  | 222.8 | 3.8 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 43 | Shajadbi Biradar    | 84 | F | 14180 | 116 | 27  | 0.6 | 189 | 92  | 48 | 122.6 | 18.4  | 4.5 | Rt. Hemiparesis | Normal       | Absent  |
| 44 | Basavaraj Jogur     | 63 | M | 7630  | 171 | 23  | 0.8 | 155 | 60  | 36 | 107   | 12    | 4.8 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 45 | Chandrawwa Golmvar  | 60 | F | 14250 | 109 | 34  | 0.8 | 224 | 105 | 62 | 141   | 21    | 5.1 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 46 | Kushaba Nalavade    | 85 | M | 15320 | 71  | 57  | 2.1 | 115 | 78  | 43 | 56.4  | 15.6  | 6.4 | Lt. Hemiparesis | Normal       | Absent  |
| 47 | Shankar Danapai     | 60 | M | 15200 | 110 | 56  | 2.3 | 240 | 119 | 42 | 174.2 | 23.8  | 4.3 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 48 | Mallappa Hittinalli | 70 | M | 6320  | 173 | 20  | 0.7 | 120 | 106 | 26 | 72.8  | 21.2  | 3.0 | Rt. Hemiparesis | Dyslipidemia | Absent  |

|    |                     |    |   |       |     |    |     |     |     |    |       |      |     |                 |              |         |
|----|---------------------|----|---|-------|-----|----|-----|-----|-----|----|-------|------|-----|-----------------|--------------|---------|
| 49 | Shrimant Padanad    | 53 | M | 9080  | 78  | 36 | 0.8 | 205 | 186 | 24 | 143.8 | 37.2 | 4.1 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 50 | Honappa Kumbar      | 70 | M | 10900 | 114 | 38 | 0.7 | 253 | 168 | 37 | 182.4 | 33.6 | 4.5 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 51 | Shivaraya Biradar   | 72 | M | 11100 | 92  | 94 | 2.6 | 135 | 121 | 20 | 90.8  | 24.2 | 7.7 | Lt. Hemiparesis | Dyslipidemia | Present |
| 52 | Sangamma Boragi     | 60 | F | 21210 | 125 | 93 | 2.8 | 131 | 105 | 35 | 75    | 21   | 8.0 | Lt. Hemiparesis | Dyslipidemia | Present |
| 53 | Basanna Hadapad     | 53 | M | 7710  | 163 | 72 | 4.0 | 155 | 112 | 57 | 75.6  | 22.4 | 6.4 | Lt. Hemiparesis | Normal       | Absent  |
| 54 | Veerendra Patil     | 42 | M | 9500  | 122 | 15 | 0.6 | 244 | 369 | 30 | 140.2 | 73.8 | 6.5 | Rt. Hemiparesis | Dyslipidemia | Absent  |
| 55 | Layawwa Ganti       | 50 | F | 10120 | 252 | 30 | 0.9 | 193 | 172 | 31 | 127.6 | 34.4 | 5.0 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 56 | Anappa Bajantri     | 70 | M | 15350 | 101 | 39 | 1.4 | 181 | 125 | 30 | 130.8 | 24.2 | 7.4 | Rt. Hemiparesis | Dyslipidemia | Present |
| 57 | Rayappa Masabinal   | 65 | M | 15990 | 212 | 28 | 1.1 | 323 | 179 | 28 | 259.2 | 35.8 | 7.8 | Rt. Hemiparesis | Dyslipidemia | Present |
| 58 | Amarayya Hiremath   | 85 | M | 21760 | 102 | 86 | 1.1 | 131 | 93  | 28 | 84.4  | 18.6 | 4.6 | Lt. Hemiparesis | Dyslipidemia | Absent  |
| 59 | Sangappa Sajjan     | 76 | M | 9040  | 113 | 25 | 1.0 | 141 | 66  | 45 | 83    | 13   | 3.1 | Lt. Hemiparesis | Normal       | Absent  |
| 60 | Gumsidawwa Bolegavi | 70 | F | 12260 | 95  | 31 | 1.0 | 323 | 179 | 28 | 259.2 | 35.8 | 7.3 | Lt. Hemiparesis | Dyslipidemia | Present |